



SABA Publishing

# Existence Theory and Stability Analysis to a Class of Hybrid Differential Equations using Conformable Fractal Fractional Derivative

SHAFIULLAH

Department of Mathematics, University of Malakand, Dir(L), Khyber Pakhtunkhwa, Pakistan

• Received: 02 April 2024

• Accepted: 20 May 2024

• Published Online: 30 May 2024

## Abstract

This study uses conformable fractal-fractional derivatives (CFFD) to investigate a class of hybrid differential equations (HDEs). Fixed point theory, in particular Krasnoselskii's fixed point theorem, is implemented on the said problem to establish the conditions for which there exists at least one solution. Also, stability results of Ulam-Hyers (U-H) and U-H Rassias types are derived. At the end of the paper, we added examples for justification and strengthening of our derived results.

Keywords: Conformable calculus, Hybrid problem, Ulam-Hyers stability.

2010 MSC: put your Mathematics Subject Classification 2010 (MSC): 26A33;34A08.

## 1. Introduction

Boundary value problems (BVPs) are important for characterizing a wide range of real-world engineering and physical research challenges. Much research has been done in the field of BVPs, which correspond to both ordinary and fractional order differential equations. Using functional analytic tools for qualitative theory, researchers have examined many BVPs. We refer [1], [2] and references there in for more details about BVPs. In [3], authors accumulated various BVPs concerned to real world problems of mathematical physics. Applications for BVPs with integral boundary conditions can be found in a wide range of fields, including population dynamics, chemical engineering, thermo-elasticity, blood flow issues, physical systems, thermo-dynamics, and other dynamics (see [4]). In recent decades, FDEs have gained much attention due to their wide use in these branches of science. Being the generalization of classical derivatives, the importance of fractional calculus is started at the same time as ordinary calculus were [5]. Using such types of fractional order derivatives have numerous benefits [6] Various definitions were defined by some researchers but the prominent role is of Riemann-Liouville and Caputo arbitrary order [7, 8].

\*Corresponding author: [sangotaswat@gmail.com](mailto:sangotaswat@gmail.com)

Recently, Atangana in [9] introduced the fractal-fractional derivatives (FFDs). Moreover, a relationship established between fractional and fractal calculus. However the application of Atangana Baleneu Fractal derivatives in neuroscience has been considered [10]. The qualitative theory of fraction DE in CFD, s very recent. Using fixed point results, for developing sufficient conditions we must see [11]. For the study of existence and stability of CFD one can see [12]. For the justification of our results, Ulam-Hyers (U-H) and U-H Rassias (U-H-R) theory as tools are helpful. The various applications to the real world problems of the CFDs [13] can strengthen our problem. The first ever used of chain rule by Khalil introduce the definition known as confirmable fractional order derivative (CFD). The properties of CFD is much similar with integer order [14]. The two operators Confirmable Derivatives and Fractal Derivatives have prominent role in theory and applications [15]. Researchers have investigated various results for different problems of fractional calculus, we refer to [16, 17, 18, 19]. Combining the two operators confirmable and fractal fractional derivatives were used recently for different problems. See some details for fractals fractional derivatives, we refer to [20, 21]. For physical applications, we refer to [22, 23].

Hybrid problems and systems of FDEs have been analyzed by researchers using different tools of mathematical analysis. In this regards, the significant contributions are cited as [24, 25, 26, 27].

We discuss our following single hybrid problem using confirmable fractional derivative. We consider the following BVP of CFFDE with integral boundary condition given by

$$\begin{cases} {}^{\text{CFF}}D^{\alpha,\beta}[\mu(t) - f(t, \mu(t))] = g(t, \mu(t)), t \in [0, b] = \mathcal{J}, \\ \mu(0) = \int_0^b h(\mu(s)) ds, \end{cases} \quad (1.1)$$

where  $f, g : \mathcal{J} \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions.

Adopting the renowned fixed point theorems of Banach and Krasnoselkii [27], adequate criteria are formulated for the uniqueness and existence of a solution. It is important to highlight that fixed point theory is a useful instrument for examining a range of issues related to the qualitative analysis. Through applications, the stability problem has attracted significant attention in a number of research domain [28], [29], and [30]. Furthermore, stability has recently been extensively investigated as a useful tool for various problems of U-H type. Recently, stability analysis of multiple types by employing various fractional differential operators has been explored by authors in [31] and [32] utilizing the U-H idea. Considering the importance of the aforementioned stability, we also look at some results for U-H and generalized U-H stabilities for our topic under consideration. Examples are given to demonstrate the results of this study.

## 2. Preliminaries

Here, we accumulate some definitions and results regarding conformable fractal fractional derivatives required in our analysis.

**Definition 2.1.** [9] The conformable derivative of a function  $f(t)$  defined on the interval  $[0, \infty)$  and for all  $t > 0$ , with order  $\alpha \in (0, 1]$ , is defined as follows:

$${}_0^{\text{C}}D_t^\alpha f(t) = \lim_{\xi \rightarrow 0} \frac{f(t + \xi t^{1-\alpha}) - f(t)}{\xi}$$

Importantly, when the function  $f$  is differentiable, we have:

$${}^C D_t^\alpha f(t) = \lim_{\xi \rightarrow 0} \frac{f(t + \xi t^{1-\alpha}) - f(t)}{\xi} = t^{1-\alpha} f'(t).$$

**Definition 2.2.** [9] The fractal derivative of a function  $f$  with order  $\alpha$  is given by:

$${}^F D_t^\alpha f(t) = \lim_{t \rightarrow t_1} \frac{f(t) - f(t_1)}{t^\alpha - t_1^\alpha}.$$

The most general case is given as:

$${}^F D_t^{\alpha, \beta} f(t) = \lim_{t \rightarrow t_1} \frac{f^\beta(t) - f^\beta(t_1)}{t^\alpha - t_1^\alpha}, \quad \alpha, \beta > 0.$$

For differentiable mapping  $f$ , one has

$$\begin{aligned} {}^F D_t^\alpha f(t) &= \lim_{t \rightarrow t_1} \frac{f(t) - f(t_1)}{t^\alpha - t_1^\alpha} = \frac{t^{1-\alpha}}{\alpha} f'(t) \\ &= \frac{1}{\alpha} {}^C D_t^\alpha f(t). \end{aligned}$$

Related integration is described as follows:

$${}^F I_t^\alpha f(t) = \alpha \int_0^t t^{\alpha-1} f(t) dt.$$

**Lemma 2.3.** [21] Let  $h \in L \in L[0, T]$ , then the solution of

$$\begin{cases} {}^{CFD} D^{\alpha, \beta} \mu(t) = h(t), & t \in \mathcal{J}, \\ \mu(0) = \mu_0 \end{cases}$$

is given by

$$\mu(t) = \mu_0 + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} h(s) ds.$$

### 3. Main Results

In this section, we present our main results.

#### 3.1. Integral representation of Problem (2.1)

The CFFDE BVP (2.1) is equivalent to the following integral equation

$$\mu(t) = \mu(0) + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds + f(t, \mu(t)), \quad t \in \mathcal{J}. \quad (3.1)$$

Let us define a Banach space say  $X = C(\mathcal{J})$  with norm defined by

$$\|\mu\| = \max_{t \in [a, b]} |\mu(t)|.$$

Now, we present some hypothesis which are helpful in building our main existence results.

(H<sub>1</sub>) There exists constants  $L_g, L_f > 0$  and  $\mu, \bar{\mu} \in X$  such that

$$\begin{aligned} |f(t, \mu(t)) - f(t, \bar{\mu}(t))| &\leq L_f |\mu - \bar{\mu}| \\ \text{and } |g(t, \mu(t)) - g(t, \bar{\mu}(t))| &\leq L_g |\mu - \bar{\mu}|. \end{aligned}$$

(H<sub>2</sub>) There exists constants  $C_g, D_g > 0$  such that

$$|g(t, \mu(t))| \leq C_g |\mu(t)| + D_g.$$

(H<sub>3</sub>) For  $\mu, \bar{\mu}$  there exists constants  $C_h > 0$  such that

$$|h(\mu) - h(\bar{\mu})| \leq C_h |\mu - \bar{\mu}|.$$

**Theorem 3.1.** *The problem (2.1) has a unique solution, if*

$$\Omega_{\alpha, \beta, L_f, L_g, C_h} = C_h + \beta L_g B(\alpha, \beta) b^{\alpha + \beta - 1} + L_f < 1. \quad (3.2)$$

*Proof.* Let  $N : X \rightarrow X$  be the operator defined by

$$N[\mu(t)] = \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds + f(t, \mu(t)).$$

Let  $\mu, \bar{\mu} \in X$ , then

$$\begin{aligned} \|N(\mu) - N(\bar{\mu})\| &= \max_{t \in [0, b]} \left| \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds + f(t, \mu(t)) \right. \\ &\quad \left. - \int_0^b h(\bar{\mu}(s)) ds - \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \bar{\mu}(s)) ds - f(t, \bar{\mu}(t)) \right| \\ &\leq \int_0^b |h(\mu(s)) - h(\bar{\mu}(s))| ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} |g(s, \mu(s)) - g(s, \bar{\mu}(s))| ds \\ &\quad + |f(t, \mu(t)) - f(t, \bar{\mu}(t))| \\ &\leq C_h \|\mu - \bar{\mu}\| b + L_g \|\mu - \bar{\mu}\| \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} ds + L_f \|\mu - \bar{\mu}\| \\ &\leq \Omega_{\alpha, \beta, L_f, L_g, C_h} \|\mu - \bar{\mu}\|. \end{aligned}$$

Hence, by inequality (3.2) operator  $N$  is a contraction. So by Banach contraction principle, the problem (2.1) has a unique solution.  $\square$

**Theorem 3.2.** *Under the hypothesis (H<sub>1</sub> – H<sub>3</sub>) the problem (2.1) has at least one solution.*

*Proof.* Let defined the space  $E = \{\mu \in X : \|\mu\| \leq \rho\}$  and taking two operators  $P, Q : X \rightarrow X$  defined by

$$\begin{aligned} Q\mu(t) &= f(t, \mu(t)), \\ P\mu(t) &= \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds. \end{aligned}$$

Then for  $\mu, \bar{\mu} \in X$ , one has

$$\begin{aligned}\|Q\mu - Q\bar{\mu}\| &= \max_{t \in \mathcal{J}} |Q\mu(t) - Q\bar{\mu}(t)| \\ &= \max_{t \in \mathcal{J}} |f(t, \mu(t)) - f(t, \bar{\mu}(t))| \\ &\leq L_f \|\mu - \bar{\mu}\|,\end{aligned}$$

then the operator  $Q$  is a contraction. Now, we show that the operator  $P$  is bounded and continuous. Let  $\mu \in E$ , then

$$\begin{aligned}\|\mu\| &= \max_{t \in [0,1]} \left| \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds \right| \\ &\leq [\rho C_h + C_0 + \beta B(\alpha, \beta) b^{\alpha+\beta-1} (\rho C_g + D_g)],\end{aligned}$$

where  $C_0 = \max_{t \in [0,1]} \int_0^b h(0) ds$ . Hence

$$\|\mu\| \leq (C_0 + \beta B(\alpha, \beta) b^{\alpha+\beta-1} C_g) \rho + C_0 + \beta B(\alpha, \beta) b^{\alpha+\beta-1} D_g \leq \rho, \quad (4)$$

where

$$\rho \geq \frac{C_g + \beta B(\alpha, \beta) b^{\alpha+\beta-1} D_g}{1 - (C_0 + \beta B(\alpha, \beta) b^{\alpha+\beta-1} C_g)}.$$

Hence  $\|\mu\| \leq \rho \Rightarrow \mu \in E$  is Bounded. In the same way, we have  $\|P\mu\| \leq \rho$  which mean  $P(E) \leq E$ .

If  $t_1 < t_2$ , then

$$\begin{aligned}|P\mu(t_2) - P\mu(t_1)| &= \left| \int_0^{t_2} \beta s^{\beta-1} (t_2 - s)^{\alpha-1} g(s, \mu(s)) ds - \int_0^{t_1} \beta s^{\beta-1} (t_1 - s)^{\alpha-1} g(s, \mu(s)) ds \right| \\ &= \left| \int_0^{t_1} [(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}] \beta s^{\beta-1} g(s, \mu(s)) ds \right. \\ &\quad \left. + \int_{t_1}^{t_2} [t_2 - s]^{\alpha-1} \beta s^{\beta-1} g(s, \mu(s)) ds \right| \\ &\leq \int_0^{t_1} (|(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}|) \beta s^{\beta-1} |g(s, \mu(s))| ds \\ &\quad + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} \beta s^{\beta-1} |g(s, \mu(s))| ds \\ &\leq \int_0^{t_1} [(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}] \beta s^{\beta-1} [C_g \|\mu\| + D_g] ds \\ &\quad + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} \beta s^{\beta-1} [C_g \|\mu\| + D_g] ds \\ &= (C_g \rho + D_g) \left[ \int_0^{t_2} \beta s^{\beta-1} (t_2 - s)^{\alpha-1} ds + \int_0^{t_1} \beta s^{\beta-1} (t_1 - s)^{\alpha-1} ds \right] \\ &= \beta (C_g \rho + D_g) B(\alpha, \beta) [t_2^{\alpha+\beta} - t_1^{\alpha+\beta}].\end{aligned}$$

Now as  $t_1 \rightarrow t_2$ , we see that right side gives us zero. Thus  $\|P\mu(t_2) - P\mu(t_1)\| \rightarrow 0$  as  $t_1 \rightarrow t_2$ . Hence  $P$  is equi-continuous. Thus by Arzelá-Ascoli theorem the operator  $P$  is compact. Consequently, it follows that problem (2.1) has at least one solution.  $\square$

#### 4. Stability Theory

Let us define a function  $\pi : \mathcal{J} \rightarrow \mathbb{R}$  independent of  $\mu$ , such that for any  $\epsilon > 0$ ,

*Remark 4.1.* (i)

$$|\pi(t)| \leq \epsilon.$$

(ii)

$$\begin{cases} {}^{\text{CFF}}\mathcal{D}^{\alpha,\beta}[\mu(t) - f(t, \mu(t))] = g(t, \mu(t)) + \pi(t), \\ \mu(0) = \int_0^b h(\mu(s)) ds, \end{cases} \quad (4.1)$$

where  $f, g : \mathcal{J} \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous functions.

The solution of (4.1) is given by

$$\mu(t) = f(t, \mu(t)) + \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{(\alpha-1)} \pi(s) ds. \quad (4.2)$$

In view of Theorem 3.1, and using Remark 4.1, one has from (4.2)

$$\begin{aligned} \mu(t) &= N(\mu(t)) + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} \pi(s) ds \\ |\mu(t) - N(\mu(t))| &\leq \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} |\pi(s)| ds \\ |\mu(t) - N(\mu(t))| &\leq b^{\alpha+\beta-1} \beta B(\alpha, \beta) \epsilon \\ \|\mu(t) - N(\mu(t))\| &\leq \Delta \epsilon, \end{aligned}$$

where

$$\Delta = b^{\alpha+\beta-1} \beta B(\alpha, \beta).$$

**Theorem 4.2.** *The solution of (2.1) is U-H stable and consequently generalized U-H stable if the condition  $\Omega_{\alpha,\beta,L_f,L_g,C_h} < 1$  holds.*

*Proof.* Consider  $\mu$  be any solution of (2.1) and  $\bar{\mu}$  be a unique solution of (2.1), then taking

$$\begin{aligned} \|\mu - \bar{\mu}\| &= \max_{t \in \mathcal{J}} |\mu(t) - \bar{\mu}(t)| \\ &\leq \max_{t \in \mathcal{J}} |\mu(t) - N\mu(t)| + \max_{t \in \mathcal{J}} |N\mu(t) - \bar{\mu}(t)|. \end{aligned}$$

Using Theorem 3.1, we have

$$\|\mu - \bar{\mu}\| \leq \Delta \epsilon + \Omega_{\alpha,\beta,L_f,L_g,C_h} \|\mu - \bar{\mu}\|.$$

After, rearrangement one has from above relation

$$\|\mu - \bar{\mu}\| \leq \frac{\Delta}{1 - \delta_{\alpha,\beta}} \epsilon.$$

Hence the solution of (2.1) is U-H stable.

In addition, let there exist a non-decreasing function  $\omega : \mathcal{J} \rightarrow \mathbb{R}$  such that  $\omega(0) = 0$ , then from above inequality we have

$$\|\mu - \bar{\mu}\| \leq \frac{\Delta}{1 - \delta_{\alpha, \beta}} \omega(\epsilon).$$

where  $\omega(\epsilon) = \epsilon$ , we see that the condition of generalized U-H stability is also holds.  $\square$

Consider the given remark

*Remark 4.3.* For function  $\pi : [0, b] \rightarrow \mathbb{R}$  independent of  $\mu$ , we have

$$|\pi(t)| \leq \lambda(t)\epsilon,$$

then the solution of

$${}^{\text{CFF}}\mathcal{D}^{\alpha, \beta} [\mu(t) - f(t, \mu(t))] = g(t, \mu(t)) + \pi(t) \quad (4.3)$$

$$\mu(0) = \int_0^b h(\mu(s)) ds \quad (4.4)$$

has

$$\mu(t) = f(t, \mu(t)) + \int_0^b h(\mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} g(s, \mu(s)) ds + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} \pi(s) ds.$$

*Proof.* In view of Theorem 3.1 yields

$$\mu(t) = \mathcal{N}(\mu(t)) + \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} \pi(s) ds, \quad t \in \mathcal{J}.$$

$$\begin{aligned} |\mu(t) - \mathcal{N}(\mu(t))| &\leq \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} |\pi(s)| ds \\ &\leq \int_0^t \beta s^{\beta-1} (t-s)^{\alpha-1} \lambda(s) \epsilon ds \\ &= \epsilon \int_0^b \beta s^{\beta-1} (b-s)^{\alpha-1} \lambda(s) ds \\ &= \epsilon \lambda_{\alpha, \beta, b}, \end{aligned}$$

where  $\lambda_{\alpha, \beta, b} = \int_0^b \beta s^{\beta-1} (b-s)^{\alpha-1} \lambda(s) ds$ . Hence, one has

$$|\mu(t) - \mathcal{N}(\mu(t))| \leq \epsilon \lambda_{\alpha, \beta, b}.$$

$\square$

**Theorem 4.4.** The solution of (2.1) is U-H Rassias stable if  $\Omega_{\alpha, \beta, L_f, L_g, C_h} < 1$ .

*Proof.* Consider  $\mu, \bar{\mu} \in X$ , then using Theorem 3.1, we have

$$\begin{aligned} \|\mu - \bar{\mu}\| &= \max_{t \in \mathcal{J}} |\mu(t) - \bar{\mu}(t)| \\ &\leq \max_{t \in \mathcal{J}} |\mu(t) - N\mu(t)| + \max_{t \in \mathcal{J}} |N\mu(t) - \bar{\mu}(t)| \\ &\leq \epsilon \lambda_{\alpha, \beta, b} + \Omega_{\alpha, \beta, L_f, L_g, C_h} \|\mu - \bar{\mu}\| \\ \|\mu - \bar{\mu}\| &\leq \frac{\lambda_{\alpha, \beta, b}}{1 - \Omega_{\alpha, \beta, L_f, L_g, C_h}} \epsilon. \end{aligned}$$

Hence the solution of (2.1) is U-H Rassias stable.

Further there exist a solution  $\psi : [0, b] \rightarrow \mathbb{R}$  non-decreasing function such that  $\psi(\epsilon) = \epsilon$ , then

$$\|\mu - \bar{\mu}\| \leq \frac{\lambda_{\alpha, \beta, b}}{1 - \Omega_{\alpha, \beta, L_f, L_g, C_h}} \psi(\epsilon).$$

which implies that the solution of considered problem is generalized U-H Rassias stable.  $\square$

## 5. Application

Here, we present examples regarding illustration of essential main results.

**Example 5.1.** Consider

$$\begin{cases} {}^{\text{CFF}}D^{\frac{1}{2}, \frac{1}{5}}[\mu(t) - f(t, \mu(t))] = g(t, \mu(t)), & t \in [0, 1], \\ \mu(0) = \int_0^1 h(\mu(s)) ds, \end{cases} \quad (5.1)$$

where

$$f(t, \mu(t)) = \frac{|\mu(t)| + 1}{t^2 + 30}, \quad g(t, \mu(t)) = \frac{|\mu(t)| + 2}{t^2 + 20} \quad \text{and} \quad h(\mu(t)) = \frac{e^{-|\mu(t)|}}{t^2 + 100}.$$

Clearly

$$|f(t, \mu(t)) - f(t, \bar{\mu}(t))| \leq \frac{1}{30} |\mu - \bar{\mu}| \quad \text{and} \quad |g(t, \mu(t)) - g(t, \bar{\mu}(t))| \leq \frac{1}{20} |\mu - \bar{\mu}|.$$

Also

$$|g(t, \mu(t))| \leq \frac{1}{20} |\mu(t)| + \frac{1}{10} \quad \text{and} \quad |h(\mu(t)) - h(\bar{\mu}(t))| \leq \frac{1}{100} |\mu - \bar{\mu}|.$$

So, we have

$$\alpha = \frac{1}{2}, \quad \beta = \frac{1}{5}, \quad b = 1, \quad L_f = \frac{1}{30}, \quad L_g = \frac{1}{20} \quad \text{and} \quad C_h = \frac{1}{100}.$$

Obviously hypothesis  $(H_1 - H_3)$  holds. Thus by Theorem 3.2, problem (5.1) has at least one solution. And

$$\Omega_{\alpha, \beta, L_f, L_g, C_h} = C_h + \beta L_g B(\alpha, \beta) b^{\alpha + \beta - 1} + L_f \approx 0.10602 < 1.$$

Hence, Theorem 3.1 implies that problem (5.1) has a unique solution. Also, assumptions of Theorem 4.2 holds, consequently the solution of (5.1) is U-H stable. In the same way, we can deduce the other kinds of U-H stability also.

**Example 5.2.** Consider

$$\begin{cases} {}^{\text{CFF}}D^{\frac{1}{3}, \frac{1}{2}}[\mu(t) - f(t, \mu(t))] = g(t, \mu(t)), & t \in [0, 1] \\ \mu(0) = \int_0^1 h(\mu(s)) ds, \end{cases} \quad (5.2)$$

where

$$f(t, \mu(t)) = \frac{\sin |\mu(t)| + 10}{t^2 + 50}, \quad g(t, \mu(t)) = \frac{\cos |\mu(t)| + 3}{t^2 + 30} \quad \text{and} \quad h(\mu(t)) = \frac{|\mu(t)|}{t^3 + 200}.$$

Clearly

$$|f(t, \mu(t)) - f(t, \bar{\mu}(t))| \leq \frac{1}{50} |\mu - \bar{\mu}| \quad \text{and} \quad |g(t, \mu(t)) - g(t, \bar{\mu}(t))| \leq \frac{1}{30} |\mu - \bar{\mu}|.$$

Also

$$|g(t, \mu(t))| \leq \frac{1}{30} |\mu(t)| + \frac{1}{10} \quad \text{and} \quad |h(\mu(t)) - h(\bar{\mu}(t))| \leq \frac{1}{200} |\mu - \bar{\mu}|.$$

So, we have

$$\alpha = \frac{1}{3}, \quad \beta = \frac{1}{2}, \quad b = 1, \quad L_f = \frac{1}{50}, \quad L_g = \frac{1}{30} \quad \text{and} \quad C_h = \frac{1}{200}.$$

Obviously hypothesis (H<sub>1</sub> – H<sub>3</sub>) holds. Thus by Theorem 3.2, problem (5.2) has at least one solution. And

$$\Omega_{\alpha, \beta, L_f, L_g, C_h} = C_h + \beta L_g B(\alpha, \beta) b^{\alpha + \beta - 1} + L_f \approx 0.05833 < 1.$$

Hence, Theorem 3.1 implies that problem (5.2) has a unique solution. Also Theorem 4.2 holds, consequently the solution to (5.2) is U-H stable. In the same way, we can deduce the other kinds of U-H stability also.

## 6. Conclusion

This work was devoted to investigate a class of HDEs with CFFD. We have deduced sufficient results for the existence theory and different kinds of U-H stability using fixed point theory and functional analysis tools. Some pertinent examples were given to demonstrate the established results. The CFFD is a powerful tool and can be used as another alternative in investigation of various dynamical problems. We concluded that the presented analysis will provide basis for deducing new results for more general nonlinear problem using CFFD. Therefore, in the future for useful results, various problems can be examined through this type of analysis.

## References

- [1] Powers, David L. (2014). Boundary value problems. Elsevier, Academic Press, New York. <https://shop.elsevier.com/books/boundary-value-problems/powers/978-0-12-374719-8>
- [2] Flyer F, and Fokas, A.S, (2008). A Hybrid Analytical-Numerical Method for Solving Evolution Partial Differential Equations. I: The Half-Line, Proc. R. Soc. London Ser. A Math Phys. Eng. Sci. **464**, 1823-1849. <https://doi.org/10.1098/rspa.2008.0041>

- [3] Ladyzhenskaya O.A. (2013). *The Boundary Value Problems of Mathematical Physics* (Vol. 49). Springer Science & Business Media, New York. <https://link.springer.com/book/10.1007/978-1-4757-4317-3>
- [4] Ahmad, B., & Nieto, J. J. (2009). Existence results for nonlinear boundary value problems of fractional integrodifferential equations with integral boundary conditions. *Boundary value problems*, **2009**, 1-11. <https://doi.org/10.1155/2009/708576>
- [5] Salah, E. Y., Sontakke, B., Abdo, M. S., Shatanawi, W., Abodayeh, K., & Albalwi, M. D. (2024). Conformable Fractional-Order Modeling and Analysis of HIV/AIDS Transmission Dynamics. *International Journal of Differential Equations*, **2024**, Article ID 1958622, 16 pages. <https://doi.org/10.1155/2024/1958622>.
- [6] Shah, K., and Abdeljawad, T. (2023). On complex fractal-fractional order mathematical modeling of CO<sub>2</sub> emanations from energy sector. *Physica Scripta*, **99**(1), 015226. DOI [10.1088/1402-4896/ad1286](https://doi.org/10.1088/1402-4896/ad1286)
- [7] Sher, M., Shah, K., Sarwar, M., Alqudah, M. A., & Abdeljawad, T. (2023). Mathematical analysis of fractional order alcoholism model. *Alexandria Engineering Journal*, **78**, 281-291. <https://doi.org/10.1016/j.aej.2023.07.010>
- [8] Suwan, I., Abdo, M., Abdeljawad, T., Mater, M., Boutiara, A., & Almalahi, M. (2021, October). Existence theorems for  $\Psi$ -fractional hybrid systems with periodic boundary conditions. *AIMS Mathematics*, **7**(1): 171-186. <https://doi.org/10.3934/math.2022010>
- [9] Atangana, A. (2017). Fractal-fractional differentiation and integration: connecting fractal calculus and fractional calculus to predict complex system. *Chaos, Solitons & Fractals*, **102**, 396-406. <https://doi.org/10.1016/j.chaos.2017.04.027>
- [10] Goufo, E. F. D., Mbehou, M., & Pene, M. M. K. (2018). A peculiar application of Atangana-Baleanu fractional derivative in neuroscience: Chaotic burst dynamics. *Chaos, Solitons & Fractals*, **115**, 170-176. <https://doi.org/10.1016/j.chaos.2018.08.003>
- [11] Lin, W. (2007). Global existence theory and chaos control of fractional differential equations. *Journal of Mathematical Analysis and Applications*, **332**(1), 709-726. <https://doi.org/10.1016/j.jmaa.2006.10.040>
- [12] Sher, M., Khan, A., Shah, K., & Abdeljawad, T. (2023). Existence and stability theory of pantograph conformable fractional differential problem. *Thermal Science*, **27**(Spec. issue 1), 237-244. <https://doi.org/10.2298/TSCI23S1237S>
- [13] Zhou, HW., Yang, S., & Zhang, SQ. (2018). Conformable derivative approach to anomalous diffusion. *Physica A: Statistical Mechanics and its Applications*, **491**, 1001-1013. <https://doi.org/10.1016/j.physa.2017.09.101>
- [14] Khalil, R., Al Horani, M., Yousef, A., & Sababheh, M. (2014). A new definition of fractional derivative. *Journal of Computational and Applied Mathematics*, **264**, 65-70. <https://doi.org/10.1016/j.cam.2014.01.002>
- [15] Alshammari, S., Alshammari, M., & Abdo, M. S. (2023). Nonlocal Hybrid Integro-Differential Equations Involving Atangana-Baleanu Fractional Operators. *Journal of Mathematics*, **2023**. <https://doi.org/10.1155/2023/5891342>
- [16] Shah, K., Arfan, M., Ullah, A., Al-Mdallal, Q., Ansari, K. J., & Abdeljawad, T. (2022). Computational study on the dynamics of fractional order differential equations with applications. *Chaos, Solitons & Fractals*, **157**, 111955. <https://doi.org/10.1016/j.chaos.2022.111955>
- [17] Dhage, BC., Dhage, SB., & Buvaneswari, K. (2019). Existence of mild solutions of nonlinear boundary value problems of coupled hybrid fractional integro differential equations. *J. Fract. Calc. Appl*, **10**(2), 191-206. <http://math-frac.org/Journals/JFCA/>
- [18] Luca, R. (2023). Advances in boundary value problems for fractional differential equations, *Fractal and Fractional*, **7**(5), 406. <https://doi.org/10.3390/fractalfract7050406>
- [19] Shah, K., & Abdeljawad, T. (2024). Study of radioactive decay process of uranium atoms via fractals-fractional analysis. *South African Journal of Chemical Engineering*, **48**, 63-70. <https://doi.org/10.1016/j.sajce.2024.01.003>
- [20] Tatom, F. B. (1995). The relationship between fractional calculus and fractals. *Fractals*, **3**(01), 217-229. <https://doi.org/10.1142/S0218348X95000175>
- [21] Atangana, A., & Khan, MA. (2019). Validity of fractal derivative to capturing chaotic attractors. *Chaos, Solitons & Fractals*, **126**, 50-59. <https://doi.org/10.1016/j.chaos.2019.06.002>
- [22] Machado, J.A. & Labora, DS. (2020). Fractional Fractals, *Fract. Calc. Appl. Anal.*, **23**, 1329-1348. <https://doi.org/10.1515/fca-2020-0066>
- [23] Chen, W., Sun, H., & Li, X. (2022). Fractal and Fractional Calculus. In *Fractional Derivative*

- Modeling in Mechanics and Engineering (pp. 83-114). Singapore: Springer Nature, Singapore. <https://link.springer.com/book/10.1007/978-981-16-8802-7>
- [24] Gul, S., Khan, RA., Khan, H., George, R., Etemad, S., & Rezapour, S. (2022). Analysis on a coupled system of two sequential hybrid BVPs with numerical simulations to a model of typhoid treatment. *Alexandria Engineering Journal*, **61**(12), 10085-10098. <https://doi.org/10.1016/j.chaos.2023.113901>
- [25] Khan, H., Ahmed, S., Alzabut, J., & Azar, A. T. (2023). A generalized coupled system of fractional differential equations with application to finite time sliding mode control for Leukemia therapy. *Chaos, Solitons & Fractals*, **174**, 113901. <https://doi.org/10.1016/j.chaos.2023.113901>
- [26] Alkhazzan, A., Al-Sadi, W., Wattanakejorn, V., Khan, H., Sitthiwirattam, T., Etemad, S., & Rezapour, S. (2022). A new study on the existence and stability to a system of coupled higher-order nonlinear BVP of hybrid FDEs under the p-Laplacian operator. *AIMS Math.*, **7**(8), 14187-14207. <http://dx.doi.org/2010.3934/math.2022782>
- [27] Latrach, K., Taoudi, MA., & Zeghal, A. (2006). Some fixed point theorems of the Schauder and the Krasnosel'skii type and application to nonlinear transport equations. *Journal of Differential Equations*, **221**(1), 256-271. <https://doi.org/10.1016/j.jde.2005.04.010>
- [28] Ren, Y., Cheng, X., & Sakthivel, R. (2014). Impulsive neutral stochastic functional integro-differential equations with infinite delay driven by fBm. *Applied Mathematics and Computation*, **247**, 205-212. <https://doi.org/10.1016/j.amc.2014.08.095>
- [29] Benzahi, A., Abada, N., Arar, N., Idris, SA., Abdo, MS., & Shatanawi, W. (2024). Caputo-Fabrizio type fractional differential equations with non-instantaneous impulses: Existence and stability results. *Alexandria Engineering Journal*, **87**, 186-200. <https://doi.org/10.1016/j.aej.2023.12.036>
- [30] Dhage, B. C. (1999). Fixed point theorems in ordered Banach algebras and applications. *Panamerican Mathematical Journal*, **9**, 83-102. [doi:10.1016/j.aml.2003.10.014](https://doi.org/10.1016/j.aml.2003.10.014)
- [31] Devi, A., & Kumar, A. (2022). Hyers-Ulam stability and existence of solution for hybrid fractional differential equation with p-Laplacian operator. *Chaos, Solitons & Fractals*, **156**, 111859. <https://doi.org/10.1016/j.chaos.2022.111859>
- [32] Shah, K., Shah, L., Ahmad, S., Rassias, JM., & Li, Y. (2021). Monotone iterative techniques together with Hyers-Ulam-Rassias stability. *Mathematical Methods in the Applied Sciences*, **44**(10), 8197-8214. <https://doi.org/10.1002/mma.5825>