

Fractional Multiplicative Corrected Dual-Simpson type inequalities

BADREDDINE MEFTAH ^{a,*} , MERIEM BOUCHARB ^b, NADJLA BOUTELHIG ^c

^a Department of Mathematics, University 8 may 1945 Guelma, Algeria.

^b Environmental Research Center (C.R.E), Annaba, Algeria.

^c Laboratoire de Recherches et des Etudes Economiques, Mohamed-Cherif Messaadia University, Souk-ahras, Algeria.

• Received: 20 October 2023 • Accepted: 10 November 2023 • Published Online: 27 December 2023

Abstract

This paper delves into the realm of inequalities, focusing on the corrected dual Simpson-type inequalities for fractional multiplicative integrals. Based on a new identity, we establish some new inequalities via multiplicative s -convexity. Finally, we provide some applications of the results obtained to special means in the frame of multiplicative calculus.

Keywords: Corrected dual-Simpson formula, integral inequalities, Fractional multiplicative integral, multiplicative s -convexity.

2010 MSC: 26A33, 26A51, 26D10, 26D15.

1. Introduction

Convexity is a fundamental mathematical concept with wide-ranging applications across various scientific disciplines. Its significance lies in its ability to capture the essential characteristics of numerous real-world phenomena, making it a potent tool for modeling and analysis. Convex functions, in particular, possess remarkable properties that simplify optimization, economics, and even the understanding of physical systems. In mathematical terms, a function φ is deemed convex on the interval $[a, b]$ if, for all $x, y \in [a, b]$, the subsequent inequality holds for all $t \in [0, 1]$:

$$\varphi(tx + (1-t)y) \leq t\varphi(x) + (1-t)\varphi(y).$$

One of the most consequential results associated with convexity is the Hermite-Hadamard (H-H) inequality. For a convex function φ defined on an interval $[a, b]$, the inequality is expressed as:

*Corresponding author: badrimeftah@yahoo.fr

$$\varphi\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b \varphi(x) dx \leq \frac{\varphi(a) + \varphi(b)}{2}.$$

The concept of convexity serves as the basis for the derivation of important integral inequalities like the Hermite-Hadamard inequality, which have wide-ranging applications in mathematical analysis, physics, and engineering, highlighting its significance in both theoretical and practical mathematical contexts, see [19, 25, 26].

Multiplicative calculus, originally introduced by Grosman and Katz in 1967, emerged as an alternative to classical calculus aimed at addressing issues concerning rates of change and multiplicative processes [13]. This mathematical framework, which exclusively deals with positive functions, underwent formalization in a comprehensive work by Bashirov et al. in 2008 [5]. Its significance lies in its enhanced capacity to handle phenomena involving growth, decay, and proportional relationships more effectively than traditional calculus. Over time, it has found relevance in a range of fields, including physics [29], finance [6], and biology [11], offering a fresh perspective for modeling and analysis in scenarios where conventional calculus may encounter limitations.

In the context of multiplicative calculus, the most suitable form is logarithmic convexity, often referred to as multiplicative convexity. This concept can be articulated as follows:

Definition 1.1 ([22]). A positive function φ is said to exhibit multiplicative convexity over the interval I if, for all $x, y \in I$, the subsequent inequality:

$$\varphi(tx + (1-t)y) \leq [\varphi(x)]^t [\varphi(y)]^{1-t},$$

holds true for all $t \in [0, 1]$.

In [2], Ali et al. established the Hermite-Hadamard inequality for multiplicative integrals as follows

$$\varphi\left(\frac{a+b}{2}\right) \leq \left(\int_a^b \varphi(x)^{dx}\right)^{\frac{1}{b-a}} \leq \sqrt{\varphi(a)\varphi(b)}, \quad (1.1)$$

where φ is multiplicatively convex on $[a, b]$.

In the domain of multiplicative integral inequalities, substantial research has been undertaken. In [7], the authors established The Midpoint and Trapezoid-type inequalities for multiplicatively convex derivatives. The Ostrowski and Simpson-type inequalities were provided by Ali et al. in [4]. The Maclaurin inequalities was provided in [18], and the Dual Simpson-type inequalities in [20]. These findings collectively enhance our understanding of multiplicative integral inequalities. For additional works, readers are referred to [3, 18, 33].

On the other hand, fractional calculus is a mathematical field that deals with differentiation and integration of non-integer order. It provides a powerful framework for modeling complex phenomena that are not always well described by traditional calculus [31, 34, 35]. In the context of fractional integral inequalities, significant progress has been made in understanding the properties of functions, fractional integration operators,

and the resulting inequalities. Researchers have explored various types of inequalities, including Hermite-Hadamard, Simpson, and Maclaurin inequalities, using fractional integral operators, see [27, 28].

The multiplicative Riemann-Liouville fractional integrals were introduced by Abdeljawad and Grossman in 2016 as follow:

Definition 1.2. [1] The multiplicative Riemann-Liouville fractional integral operators of order $\alpha \in \mathbb{C}$, where $\operatorname{Re}(\alpha) > 0$, are defined as follows:

$${}_a I_*^\alpha \varphi(x) = e^{(J_{a+}^\alpha (\ln \circ \varphi))(x)}, \quad a < x \quad (1.2)$$

and

$${}_b I_*^\alpha \varphi(x) = e^{(J_{b-}^\alpha (\ln \circ \varphi))(x)}, \quad x < b, \quad (1.3)$$

where J_{c+}^α and J_{c-}^α represent the classical Riemann-Liouville operators.

In [9], Budak and Özçelik, established the Hermite-Hadamard inequalities pertaining to multiplicative Riemann-Liouville fractional integrals as follow:

$$\varphi\left(\frac{a+b}{2}\right) \leq \left[{}_a I_{\frac{a+b}{2}}^\alpha \varphi(a) \frac{a+b}{2} I_*^\alpha \varphi(b) \right]^{\frac{2\alpha-1\Gamma(\alpha+1)}{(b-a)^\alpha}} \leq \sqrt{\varphi(a)\varphi(b)}, \quad (1.4)$$

where φ is multiplicatively convex function defined on the interval $[a, b]$, and $\alpha > 0$.

A generalization of the results presented by Ali et al. in [2] and Budak et al. in [9] was provided by Fu et al. in [12]. The fractional multiplicative Bullen-type inequalities is given in [8], while the Simpson-type inequalities were provided in [21]. Moreover, Peng et al. made a significant contribution to the field through their investigation into fractional multiplicative Maclaurin-type inequalities in [24]. Additional pertinent results are available in [23, 30], as well as in the referenced sources.

The corrected dual-Simpson formula for multiplicative integrals is given by:

$$\left(\int_a^b \varphi(x) dx \right)^{\frac{1}{b-a}} = \left[\left(\varphi\left(\frac{3a+b}{4}\right) \right)^{\frac{8}{15}} \left(\varphi\left(\frac{a+b}{2}\right) \right)^{\frac{-1}{15}} \left(\varphi\left(\frac{a+3b}{4}\right) \right)^{\frac{8}{15}} \right] + \mathcal{R}(\varphi),$$

where $\mathcal{R}(\varphi)$ is the approximation error, see [16].

Motivated by the above cited papers, in this work, we will discuss the corrected dual Simpson's formula via fractional multiplicative integrals. To do so, we first establish a new integral identity. On the basis of this equality, we derive some corrected dual-Simpson type inequalities for functions whose multiplicative derivatives are multiplicatively s -convex in the second sense.

We recall that a function is said to be multiplicatively s -convex in the second sense for some fixed $s \in (0, 1]$ if, for all $x, y \in I$ the following inequality holds true for all $t \in [0, 1]$.

$$\varphi(tx + (1-t)y) \leq [\varphi(x)]^{t^s} [\varphi(y)]^{(1-t)^s}.$$

2. Preliminaries

Definition 2.1 ([5]). The definition of the multiplicative derivative for a positive function φ , indicated as φ^* , is provided as follows:

$$\frac{d^*\varphi}{dx} = \varphi^*(x) = \lim_{h \rightarrow 0} \left(\frac{\varphi(x+h)}{\varphi(x)} \right)^{\frac{1}{h}}.$$

Remark 2.2. Every positive and differentiable function φ naturally possesses multiplicative differentiability, with the relationship between φ' and φ^* determined by the following connection:

$$\varphi^*(x) = e^{(\ln \varphi(x))'} = e^{\frac{\varphi'(x)}{\varphi(x)}}.$$

Proposition 2.3 ([5]). Suppose φ and ξ are two multiplicatively differentiable functions, and ψ is differentiable. Let k be an arbitrary positive constant. Then, the functions $k\varphi$, $\varphi\xi$, $\varphi + \xi$, φ/ξ , and φ^ψ exhibit multiplicative differentiability, and

- $(k\varphi)^*(x) = \varphi^*(x),$
- $(\varphi + \xi)^*(x) = \varphi^*(x)^{\frac{\varphi(x)}{\varphi(x)+\xi(x)}} \xi^*(x)^{\frac{\xi(x)}{\varphi(x)+\xi(x)}},$
- $(\varphi\xi)^*(x) = \varphi^*(x) \xi^*(x),$
- $\left(\frac{\varphi}{\xi}\right)^*(x) = \frac{\varphi^*(x)}{\xi^*(x)},$
- $(\varphi^\psi)^*(x) = \varphi^*(x)^{\psi(x)} \varphi(x)^{\psi'(x)},$
- $(\varphi \circ \psi)^*(x) = \varphi^*(\psi(x))^{\psi'(x)}.$

Definition 2.4 ([5]). The definition of the multiplicative integral for a positive function φ , is provided as follows:

$$\int_a^b (\varphi(x))^{dx} = \exp \left(\int_a^b \ln(\varphi(x)) dx \right).$$

Proposition 2.5 ([5]). If we have positive and Riemann-integrable functions φ and ξ , then both φ and ξ are considered multiplicative integrable, and the following properties are valid.

- $\int_a^b ((\varphi(x))^p)^{dx} = \left(\int_a^b (\varphi(x))^{dx} \right)^p, \quad p \in \mathbb{R},$
- $\int_a^b (\varphi(x))^{dx} = \int_a^c (\varphi(x))^{dx} \int_c^b (\varphi(x))^{dx}, \quad a < c < b,$
- $\int_a^a (\varphi(x))^{dx} = 1$ and $\int_a^b (\varphi(x))^{dx} = \left(\int_b^a (\varphi(x))^{dx} \right)^{-1},$

$$\begin{aligned} & \bullet \int_a^b (\varphi(x) \xi(x))^{dx} = \int_a^b (\varphi(x))^{dx} \int_a^b (\xi(x))^{dx}, \\ & \bullet \int_a^b \left(\frac{\varphi(x)}{\xi(x)} \right)^{dx} = \frac{\int_a^b (\varphi(x))^{dx}}{\int_a^b (\xi(x))^{dx}}. \end{aligned}$$

The multiplicative integration by parts is as follows:

Lemma 2.6 ([5]). *Take a positive multiplicative differentiable function φ defined on the interval $[a, b]$. Also, let $\psi : J \subset \mathbb{R} \rightarrow \mathbb{R}$ and $\xi : [a, b] \rightarrow \mathbb{R}$ be two separate differentiable functions. Consequently, we can state:*

$$\int_a^b \left(\varphi^*(\psi(x)) \psi'(x) \xi(x) \right)^{dx} = \frac{\varphi(\psi(b))^{\xi(b)}}{\varphi(\psi(a))^{\xi(a)}} \times \frac{1}{\int_a^b \left(\varphi(\psi(x))^{\xi'(x)} \right)^{dx}}.$$

Now, let's recall particular functions that will be used later in our inquiry.

Definition 2.7. [15] The beta function is defined for any complex numbers x, y such that $\operatorname{Re}(x) > 0$ and $\operatorname{Re}(y) > 0$ by

$$B(x, y) = \int_0^1 v^{x-1} (1-v)^{y-1} dv = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)},$$

where $\Gamma(\cdot)$ denotes the Euler gamma function.

Definition 2.8. [15] For any complex numbers x, y, z and a such that $\operatorname{Re}(z) > \operatorname{Re}(y) > 0$ and $|a| < 1$, the hypergeometric function is defined as follows

$${}_2F_1(x, y, z; y) = \frac{1}{B(y, z-y)} \int_0^1 v^{y-1} (1-v)^{z-y-1} (1-av)^{-x} dv,$$

where $B(\cdot, \cdot)$ is the beta function.

Here is an example from biology that allows us to assess the usefulness of the concepts of multiplicative differentiation and integration.

Example 2.9. In biology, a common example that can be modeled using an ordinary differential equation (ODE) involves the growth of a population of bacteria. Consider a population of bacteria in an ideal environment with sufficient resources for their growth. We can use an ODE to represent the change in the population size of bacteria over time.

One of the simplest population growth equations for bacteria is the exponential growth equation:

$$\varphi'(t) = \mathcal{R}(t) \times \varphi(t), \quad (2.1)$$

where:

- φ is the size of the bacterial population.
- φ' represents the rate of change in the bacterial population with respect to time.
- $\mathcal{R}(t)$ is the intrinsic growth rate of the bacterial population.

This equation models bacterial population growth exponentially, meaning that the population increases proportionally to its current size. However, in reality, bacterial growth is subject to limiting factors, such as the availability of finite resources, which can be accounted for using more complex growth equations like the logistic growth equation.

It is important to note that the time variable, t , is always positive and the growth rate of the bacterial population is positive.

Using multiplicative calculus, equation (2.1) can be reformulated as $e^{(\ln(\varphi(t)))'} = e^{\mathcal{R}(t)}$, or $\varphi^*(t) = e^{\mathcal{R}(t)}$, whose solution is given in the form of a multiplicative integral, as follows:

$$\varphi(t) = \lambda \int_{t_0}^t \left(e^{\mathcal{R}(t)} \right)^{dt}, \quad \text{with } \lambda = \varphi(t_0).$$

Hence, through this example, we have assessed the usefulness and connection between the concepts of multiplicative differentiation and integration and their application to differential equations.

3. Main results

In order to establish our results, we need the following lemma.

Lemma 3.1. *Assuming that φ is a positive multiplicatively differentiable function on $[a, b]$, and φ^* is multiplicatively integrable over $[a, b]$, then for $\alpha > 0$, $\rho \geq 0$ and $x \in [a, \frac{a+b}{2}]$, the following fractional multiplicative integral identity holds:*

$$\begin{aligned} & \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] (\mathcal{J}(a, b; \varphi))^{-\frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1)} \\ &= \left(\int_0^1 \left(\left(\varphi^* \left((1-t)a + t\frac{3a+b}{4} \right) \right)^{t^\alpha} \right)^{dt} \right)^{\frac{b-a}{16}} \\ & \times \left(\int_0^1 \left(\left(\varphi^* \left((1-t)\frac{3a+b}{4} + t\frac{a+b}{2} \right) \right)^{-\frac{2}{15} - (1-t)^\alpha} \right)^{dt} \right)^{\frac{b-a}{16}} \\ & \times \left(\int_0^1 \left(\left(\varphi^* \left((1-t)\frac{a+b}{2} + t\frac{a+3b}{4} \right) \right)^{t^\alpha + \frac{2}{15}} \right)^{dt} \right)^{\frac{b-a}{16}} \\ & \times \left(\int_0^1 \left(\left(\varphi^* \left((1-t)\frac{a+3b}{4} + tb \right) \right)^{-(1-t)^\alpha} \right)^{dt} \right)^{\frac{b-a}{16}}, \end{aligned}$$

where $\mathcal{J}$ is defined as follows:

$$\begin{aligned} \mathcal{J}(a, b; \varphi) = & \left(\left({}^*I_{\left(\frac{3a+b}{4}\right)}^{\alpha} \varphi \right) (a) \left(\left(\frac{a+3b}{4} \right) I_*^{\alpha} \varphi \right) (b) \right. \\ & \times \left. \left(\left(\frac{3a+b}{4} \right) I_*^{\alpha} \varphi \right) \left(\frac{a+b}{2} \right) \left({}^*I_{\left(\frac{a+3b}{4}\right)}^{\alpha} \varphi \right) \left(\frac{a+b}{2} \right) \right)^{\frac{1}{(b-a)^{\alpha-1}}}. \end{aligned} \quad (3.1)$$

Proof. Let

$$\begin{aligned} I_1 &= \left(\int_0^1 \left((\varphi^* ((1-t)a + t\frac{3a+b}{4})) t^{\alpha} \right) dt \right)^{\frac{b-a}{16}}, \\ I_2 &= \left(\int_0^1 \left((\varphi^* ((1-t)\frac{3a+b}{4} + t\frac{a+b}{2}))^{-\frac{2}{15} - (1-t)^{\alpha}} \right) dt \right)^{\frac{b-a}{16}}, \\ I_3 &= \left(\int_0^1 \left((\varphi^* ((1-t)\frac{a+b}{2} + t\frac{a+3b}{4})) t^{\alpha + \frac{2}{15}} \right) dt \right)^{\frac{b-a}{16}} \end{aligned}$$

and

$$I_4 = \left(\int_0^1 \left((\varphi^* ((1-t)\frac{a+3b}{4} + tb))^{-(1-t)^{\alpha}} \right) dt \right)^{\frac{(x-a)^2}{b-a}}.$$

Using the multiplicative integration by parts, from I_1 we have

$$\begin{aligned} I_1 &= \left(\int_0^1 \left((\varphi^* ((1-t)a + t\frac{3a+b}{4})) t^{\alpha} \right) dt \right)^{\frac{b-a}{16}} \\ &= \int_0^1 \left((\varphi^* ((1-t)a + t\frac{3a+b}{4}))^{\frac{b-a}{16} t^{\alpha}} \right) dt \\ &= (\varphi(\frac{3a+b}{4}))^{\frac{b-a}{4}} \frac{1}{\int_0^1 \left((\varphi((1-t)a + t\frac{3a+b}{4}))^{\alpha \frac{b-a}{4} t^{\alpha-1}} \right) dt} \\ &= (\varphi(\frac{3a+b}{4}))^{\frac{b-a}{4}} \frac{1}{\exp \left\{ \alpha \frac{b-a}{4} \int_0^1 \left(t^{\alpha-1} (\ln \varphi((1-t)a + t\frac{3a+b}{4})) \right) dt \right\}} \\ &= (\varphi(\frac{3a+b}{4}))^{\frac{b-a}{4}} \frac{1}{\exp \left\{ \frac{\alpha 4^{\alpha-1}}{(b-a)^{\alpha}} \int_a^{\frac{3a+b}{4}} ((y-a)^{\alpha-1} (\ln \varphi(y))) dy \right\}} \\ &= (\varphi(\frac{3a+b}{4}))^{\frac{b-a}{4}} \frac{1}{\left(\exp \left\{ \left(\frac{1}{\Gamma(\alpha)} \int_a^{\frac{3a+b}{4}} ((y-a)^{\alpha-1} (\ln \varphi(y))) dy \right) \right\} \right)^{\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}}} \end{aligned}$$

$$= \left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{b-a}{4}} \left(\left({}^* I_{\left(\frac{3a+b}{4} \right)}^{\alpha} \varphi \right) (a) \right)^{-\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}}. \quad (3.2)$$

Similarly,

$$\begin{aligned} I_2 &= \left(\int_0^1 \left(\left(\varphi^* \left((1-t) \frac{3a+b}{4} + t \frac{a+b}{2} \right) \right)^{-\frac{2}{15} - (1-t)^{\alpha}} \right)^{\frac{b-a}{16}} dt \right) \\ &= \left(\int_0^1 \left(\left(\varphi^* \left((1-t) \frac{3a+b}{4} + t \frac{a+b}{2} \right) \right)^{\frac{b-a}{16} \left(-\frac{2}{15} - (1-t)^{\alpha} \right)} \right)^{\frac{b-a}{16}} dt \right) \\ &= \frac{\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{17}{60}}}{\left(\varphi \left(\frac{a+b}{2} \right) \right)^{\frac{1}{30}}} \frac{1}{\int_0^1 \left(\left(\varphi \left((1-t) \frac{3a+b}{4} + t \frac{a+b}{2} \right) \right)^{\frac{\alpha}{4} (1-t)^{\alpha-1}} \right)^{\frac{b-a}{16}} dt} \\ &= \frac{\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{17}{60}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}}}{\exp \left\{ \frac{\alpha}{4} \int_0^1 (1-t)^{\alpha-1} \ln \left(\varphi \left((1-t) \frac{3a+b}{4} + t \frac{a+b}{2} \right) \right) dt \right\}} \\ &= \frac{\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{17}{60}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}}}{\exp \left\{ \frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}} \left(\frac{1}{\Gamma(\alpha)} \int_{\frac{3a+b}{4}}^{\frac{a+b}{2}} \left(\frac{a+b}{2} - y \right)^{\alpha-1} \ln(\varphi(y)) dy \right) \right\}} \\ &= \left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{17}{60}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}} \left(\left({}^* I_{\left(\frac{3a+b}{4} \right)}^{\alpha} \varphi \right) \left(\frac{a+b}{2} \right) \right)^{-\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}}, \quad (3.3) \end{aligned}$$

$$\begin{aligned} I_3 &= \left(\int_0^1 \left(\left(\varphi^* \left((1-t) \frac{a+b}{2} + t \frac{a+3b}{4} \right) \right)^{t^{\alpha} + \frac{2}{15}} \right)^{\frac{b-a}{16}} dt \right) \\ &= \int_0^1 \left(\left(\varphi^* \left((1-t) \frac{a+b}{2} + t \frac{a+3b}{4} \right) \right)^{\frac{b-a}{16} \left(t^{\alpha} + \frac{2}{15} \right)} \right)^{\frac{b-a}{16}} dt \\ &= \frac{\left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{17}{60}}}{\left(\varphi \left(\frac{a+b}{2} \right) \right)^{\frac{1}{30}}} \frac{1}{\int_0^1 \left(\left(\varphi \left((1-t) \frac{a+b}{2} + t \frac{a+3b}{4} \right) \right)^{\frac{\alpha}{4} t^{\alpha-1}} \right)^{\frac{b-a}{16}} dt} \\ &= \frac{\left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{17}{60}}}{\exp \left\{ \frac{\alpha}{4} \int_0^1 t^{\alpha-1} \ln \left(\varphi \left((1-t) \frac{a+b}{2} + t \frac{a+3b}{4} \right) \right) dt \right\}} \\ &= \frac{\left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{17}{60}}}{\left(\exp \left\{ \frac{1}{\Gamma(\alpha)} \int_{\frac{a+b}{2}}^{\frac{a+3b}{4}} \left(y - \frac{a+b}{2} \right)^{\alpha-1} \ln(\varphi(y)) dy \right\} \right)^{\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}}} \\ &= \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{30}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{17}{60}} \left(\left({}^* I_{\left(\frac{a+3b}{4} \right)}^{\alpha} \varphi \right) \left(\frac{a+b}{2} \right) \right)^{-\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}} \quad (3.4) \end{aligned}$$

and

$$\begin{aligned}
 I_4 &= \left(\int_0^1 \left((\varphi^* \left((1-t) \frac{a+3b}{4} + tb \right))^{-(1-t)^\alpha} \right)^{\frac{b-a}{16}} dt \right) \\
 &= \left(\int_0^1 \left((\varphi^* \left((1-t) \frac{a+3b}{4} + tb \right))^{-\frac{b-a}{16}(1-t)^\alpha} \right)^{\frac{b-a}{16}} dt \right) \\
 &= \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{1}{4}} \frac{1}{\int_0^1 \left(\varphi \left((1-t) \frac{a+3b}{4} + tb \right) \right)^{\frac{\alpha}{4}(1-t)^{\alpha-1}} dt} \\
 &= \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{1}{4}} \frac{1}{\exp \left\{ \frac{\alpha}{4} \int_0^1 (1-t)^{\alpha-1} \ln \left(\varphi \left((1-t) \frac{a+3b}{4} + tb \right) \right) dt \right\}} \\
 &= \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{1}{4}} \frac{1}{\exp \left\{ \frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1) \left(\frac{1}{\Gamma(\alpha)} \int_{\frac{a+3b}{4}}^b (b-y)^{\alpha-1} \ln(\varphi(y)) dy \right) \right\}} \\
 &= \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{1}{4}} \left(\left(\frac{a+3b}{4} \right) I_*^\alpha \varphi \right) (b) \right)^{-\frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1)}. \tag{3.5}
 \end{aligned}$$

By multiplying (3.2)-(3.5) we obtain the desired result. This completes the proof. $\square$

Theorem 3.2. Let $\varphi : [a, b] \rightarrow \mathbb{R}^+$ be an increasing and multiplicatively differentiable function on $[a, b]$. If φ^* is multiplicative s -convex on $[a, b]$, then we have

$$\begin{aligned}
 &\left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] (\mathcal{J}(a, b; \varphi))^{-\frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1)} \right| \\
 &\leq \left[(\varphi^*(a) \varphi^*(b))^{B(\alpha+1, s+1)} \varphi^* \left(\frac{a+b}{2} \right) \right]^{\left(\frac{4}{15(s+1)} + 2B(s+1, \alpha+1) \right)} \\
 &\quad \times \left(\varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{32(s+1)+2\alpha}{15(\alpha+s+1)(s+1)} \right)} \right]^{\frac{b-a}{16}}.
 \end{aligned}$$

where $\mathcal{J}$ is defined as (3.1), and B is the beta function.

Proof. From Lemma 3.1, properties of multiplicative integrals and the multiplicative s -convexity of φ^* , we have

$$\begin{aligned}
 &\left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] (\mathcal{J}(a, b; \varphi))^{-\frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1)} \right| \\
 &\leq \exp \left(\frac{b-a}{16} \int_0^1 t^\alpha \ln \left(\varphi^* \left((1-t)a + t \frac{3a+b}{4} \right) \right) dt \right) \\
 &\quad \times \exp \left(\frac{(a+b-2x)^2}{4(b-a)} \int_0^1 \left(\frac{2}{15} + (1-t)^\alpha \right) \ln \varphi^* \left((1-t) \frac{3a+b}{4} + t \frac{a+b}{2} \right) dt \right)
 \end{aligned}$$

$$\begin{aligned}
& \times \exp \left(\frac{b-a}{16} \int_0^1 \left(t^\alpha + \frac{2}{15} \right) \ln \varphi^* \left((1-t) \frac{a+b}{2} + t \frac{a+3b}{4} \right) dt \right) \\
& \times \exp \left(\frac{b-a}{16} \int_0^1 (1-t)^\alpha \ln \varphi^* \left((1-t) \frac{a+3b}{4} + tb \right) dt \right) \\
& \leq \exp \left(\frac{b-a}{16} \int_0^1 t^\alpha \left((1-t)^s \ln \varphi^*(a) + t^s \ln \varphi^* \left(\frac{3a+b}{4} \right) \right) dt \right) \\
& \times \exp \left(\frac{b-a}{16} \int_0^1 \left(\frac{2}{15} + (1-t)^\alpha \right) \left((1-t)^s \ln \varphi^* \left(\frac{3a+b}{4} \right) + t^s \ln \varphi^* \left(\frac{a+b}{2} \right) \right) dt \right) \\
& \times \exp \left(\frac{b-a}{16} \int_0^1 \left(t^\alpha + \frac{2}{15} \right) \left((1-t)^s \ln \varphi^* \left(\frac{a+b}{2} \right) + t^s \ln \varphi^* \left(\frac{a+3b}{4} \right) \right) dt \right) \\
& \times \exp \left(\frac{b-a}{16} \int_0^1 (1-t)^\alpha \left((1-t)^s \ln \varphi^* \left(\frac{a+3b}{4} \right) + t^s \ln \varphi^*(b) \right) dt \right) \\
& = \exp \left(\frac{b-a}{16} \left(\int_0^1 t^\alpha (1-t)^s dt \right) \ln \varphi^*(a) + \frac{b-a}{16} \left(\int_0^1 t^{\alpha+s} dt \right) \ln \varphi^* \left(\frac{3a+b}{4} \right) \right) \\
& \times \exp \left(\frac{b-a}{16} \left(\int_0^1 \left(\frac{2}{15} + (1-t)^\alpha \right) (1-t)^s dt \right) \ln \varphi^* \left(\frac{3a+b}{4} \right) \right. \\
& \left. + \frac{b-a}{16} \left(\int_0^1 \left(\frac{2}{15} + (1-t)^\alpha \right) t^s dt \right) \ln \varphi^* \left(\frac{a+b}{2} \right) \right) \\
& \times \exp \left(\frac{b-a}{16} \left(\int_0^1 \left(t^\alpha + \frac{2}{15} \right) (1-t)^s dt \right) \ln \varphi^* \left(\frac{a+b}{2} \right) \right. \\
& \left. + \frac{b-a}{16} \left(\int_0^1 \left(t^\alpha + \frac{2}{15} \right) t^s dt \right) \ln \varphi^* \left(\frac{a+3b}{4} \right) dt \right) \\
& \times \exp \left(\frac{b-a}{16} \left(\int_0^1 (1-t)^{\alpha+s} dt \right) \ln \varphi^* \left(\frac{a+3b}{4} \right) + \frac{b-a}{16} \left(\int_0^1 (1-t)^\alpha t^s dt \right) \ln \varphi^*(b) \right) \\
& = \left[(\varphi^*(a) \varphi^*(b))^{\mathbf{B}(\alpha+1, s+1)} \varphi^* \left(\frac{a+b}{2} \right)^{\left(\frac{4}{15(s+1)} + 2\mathbf{B}(s+1, \alpha+1) \right)} \right. \\
& \left. \times (\varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right))^{\left(\frac{32(s+1)+2\alpha}{15(\alpha+s+1)(s+1)} \right)} \right]^{\frac{b-a}{16}},
\end{aligned}$$

where we have used

$$\int_0^1 t^\alpha (1-t)^s dt = \int_0^1 (1-t)^\alpha t^s dt = B(\alpha+1, s+1), \quad (3.6)$$

$$\int_0^1 t^{\alpha+s} dt = \int_0^1 (1-t)^{\alpha+s} dt = \frac{1}{\alpha+s+1}, \quad (3.7)$$

$$\int_0^1 \left(\frac{2}{15} + (1-t)^\alpha\right) (1-t)^s dt = \int_0^1 \left(t^\alpha + \frac{2}{15}\right) t^s dt = \frac{17(s+1)+2\alpha}{15(\alpha+s+1)(s+1)} \quad (3.8)$$

and

$$\int_0^1 \left(t^\alpha + \frac{2}{15}\right) (1-t)^s dt = \int_0^1 \left(\frac{2}{15} + (1-t)^\alpha\right) t^s dt = \frac{2}{15(s+1)} + B(s+1, \alpha+1). \quad (3.9)$$

The proof is completed. $\square$

Corollary 3.3. *In Theorem 3.2, if we take $s = 1$, then we get the following fractional corrected dual-Simpson type inequality for multiplicative convex functions*

$$\begin{aligned} & \left| \left[\left(\varphi\left(\frac{3a+b}{4}\right) \right)^{\frac{8}{15}} \left(\varphi\left(\frac{a+b}{2}\right) \right)^{-\frac{1}{15}} \left(\varphi\left(\frac{a+3b}{4}\right) \right)^{\frac{8}{15}} \right] (J(a, b; \varphi))^{-\frac{4\alpha-1}{(b-a)^\alpha} \Gamma(\alpha+1)} \right| \\ & \leq \left[\left(\varphi^*(a) \varphi^*(b) \right)^{\frac{1}{(\alpha+1)(\alpha+2)}} \varphi^*\left(\frac{a+b}{2}\right)^{\left(\frac{2(\alpha+1)(\alpha+2)+30}{15(\alpha+1)(\alpha+2)}\right)} \right. \\ & \quad \left. \times \left(\varphi^*\left(\frac{3a+b}{4}\right) \varphi^*\left(\frac{a+3b}{4}\right) \right)^{\left(\frac{32+\alpha}{15(\alpha+2)}\right)} \right]^{\frac{b-a}{16}}. \end{aligned}$$

Corollary 3.4. *By setting $\alpha = 1$ in Theorem 3.2, we get*

$$\begin{aligned} & \left| \left[\left(\varphi\left(\frac{3a+b}{4}\right) \right)^{\frac{8}{15}} \left(\varphi\left(\frac{a+b}{2}\right) \right)^{-\frac{1}{15}} \left(\varphi\left(\frac{a+3b}{4}\right) \right)^{\frac{8}{15}} \right] \left(\int_a^b \varphi(x) dx \right)^{\frac{1}{a-b}} \right| \\ & \leq \left[\left(\varphi^*(a) \varphi^*(b) \right)^{\frac{1}{(s+1)(s+2)}} \varphi^*\left(\frac{a+b}{2}\right)^{\left(\frac{4(s+2)+30}{15(s+1)(s+2)}\right)} \right. \\ & \quad \left. \times \left(\varphi^*\left(\frac{3a+b}{4}\right) \varphi^*\left(\frac{a+3b}{4}\right) \right)^{\left(\frac{32(s+1)+2}{15(s+1)(s+2)}\right)} \right]^{\frac{b-a}{16}}. \end{aligned}$$

Corollary 3.5. *In Theorem 3.2, if we take $\alpha = s = 1$, then we get*

$$\begin{aligned} & \left| \left[\left(\varphi\left(\frac{3a+b}{4}\right) \right)^{\frac{8}{15}} \left(\varphi\left(\frac{a+b}{2}\right) \right)^{-\frac{1}{15}} \left(\varphi\left(\frac{a+3b}{4}\right) \right)^{\frac{8}{15}} \right] \left(\int_a^b \varphi(x) dx \right)^{\frac{1}{a-b}} \right| \\ & \leq \left[\left(\varphi^*(a) \varphi^*(b) \right)^{\frac{1}{6}} \varphi^*\left(\frac{a+b}{2}\right)^{\frac{7}{15}} \left(\varphi^*\left(\frac{3a+b}{4}\right) \varphi^*\left(\frac{a+3b}{4}\right) \right)^{\left(\frac{11}{15}\right)} \right]^{\frac{b-a}{16}}. \end{aligned}$$

Theorem 3.6. Let $\varphi : [a, b] \rightarrow \mathbb{R}^+$ be an increasing and multiplicatively differentiable over $[a, b]$. If for $q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, $(\ln \varphi^*)^q$ is s -convex on $[a, b]$, then we have

$$\begin{aligned} & \left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{\frac{-1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] (\mathcal{J}(a, b; \varphi))^{-\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}} \right| \\ & \leq \left[\left(\varphi^*(a) \varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \varphi^*(b) \right)^{\left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}}} \right. \\ & \quad \times \left. \left(\varphi^* \left(\frac{3a+b}{4} \right) \left(\varphi^* \left(\frac{a+b}{2} \right) \right)^2 \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\frac{17}{15} \right) \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}}} \right]^{\frac{b-a}{16} \frac{1}{q}}, \end{aligned}$$

where $\mathcal{J}$ is defined as (3.1), and ${}_2F_1$ is the hypergeometric function.

Proof. From Lemma 3.1, Hölder inequality and s -convexity of $(\ln \varphi^*)^q$, we have:

$$\begin{aligned} & \left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{\frac{-1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] (\mathcal{J}(a, b; \varphi))^{-\frac{4^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^{\alpha}}} \right| \\ & \leq \exp \left(\frac{b-a}{16} \left(\int_0^1 t^{\alpha p} dt \right)^{\frac{1}{p}} \left(\int_0^1 |\ln(\varphi^*((1-t)a + t \frac{3a+b}{4}))|^q dt \right)^{\frac{1}{q}} \right) \\ & \quad \times \exp \left(\frac{b-a}{16} \left(\int_0^1 ((1-t)^{\alpha} + \frac{2}{15})^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |\ln \varphi^*((1-t) \frac{3a+b}{4} + t \frac{a+b}{2})|^q dt \right)^{\frac{1}{q}} \right) \\ & \quad \times \exp \left(\frac{b-a}{16} \left(\int_0^1 (t^{\alpha} + \frac{2}{15})^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |\ln \varphi^*((1-t) \frac{a+b}{2} + t \frac{a+3b}{4})|^q dt \right)^{\frac{1}{q}} \right) \\ & \quad \times \exp \left(\frac{b-a}{16} \left(\int_0^1 (1-t)^{\alpha p} dt \right)^{\frac{1}{p}} \left(\int_0^1 |\ln \varphi^*((1-t) \frac{a+3b}{4} + tb)|^q dt \right)^{\frac{1}{q}} \right) \\ & \leq \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \right. \\ & \quad \times \left. \left(\int_0^1 \left((1-t)^s (\ln(\varphi^*(a)))^q + t^s (\ln(\varphi^*(\frac{3a+b}{4})))^q \right) dt \right)^{\frac{1}{q}} \right) \\ & \quad \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \right. \\ & \quad \times \left. \left(\int_0^1 \left((1-t)^s (\ln \varphi^*(\frac{3a+b}{4}))^q + t^s (\ln \varphi^*(\frac{a+b}{2}))^q \right) dt \right)^{\frac{1}{q}} \right) \end{aligned}$$

$$\begin{aligned}
& \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \right. \\
& \times \left. \left(\int_0^1 \left((1-t)^s \left(\ln \varphi^* \left(\frac{a+b}{2} \right) \right)^q + t^s \left(\ln \varphi^* \left(\frac{a+3b}{4} \right) \right)^q \right) dt \right)^{\frac{1}{q}} \right) \\
& \times \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\int_0^1 \left((1-t)^s \left(\ln \varphi^* \left(\frac{a+3b}{4} \right) \right)^q + t^s \left(\ln \varphi^* (b) \right)^q \right) dt \right)^{\frac{1}{q}} \right) \\
& = \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\left(\ln \left(\varphi^* (a) \right) \right)^q + \left(\ln \left(\varphi^* \left(\frac{3a+b}{4} \right) \right) \right)^q \right)^{\frac{1}{q}} \right) \\
& \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \right. \\
& \times \left. \left(\left(\ln \varphi^* \left(\frac{3a+b}{4} \right) \right)^q + \left(\ln \varphi^* \left(\frac{a+b}{2} \right) \right)^q \right)^{\frac{1}{q}} \right) \\
& \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \right. \\
& \times \left. \left(\left(\ln \varphi^* \left(\frac{a+b}{2} \right) \right)^q + \left(\ln \varphi^* \left(\frac{a+3b}{4} \right) \right)^q \right)^{\frac{1}{q}} \right) \\
& \times \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\left(\ln \varphi^* \left(\frac{a+3b}{4} \right) \right)^q + \left(\ln \varphi^* (b) \right)^q \right)^{\frac{1}{q}} \right), \tag{3.10}
\end{aligned}$$

where we have used

$$\begin{aligned}
\int_0^1 \left((1-t)^\alpha + \frac{2}{15} \right)^p dt &= \int_0^1 \left(t^\alpha + \frac{2}{15} \right)^p dt = \frac{1}{\alpha} \int_0^1 \left(u + \frac{2}{15} \right)^p u^{\frac{1}{\alpha}-1} du \\
&= \frac{1}{\alpha} \int_0^1 \left(\frac{17}{15} - x \right)^p (1-x)^{\frac{1}{\alpha}-1} dx \\
&= \left(\frac{17}{15} \right)^p \int_0^1 \left(1 - \frac{15}{17}x \right)^p (1-x)^{\frac{1}{\alpha}-1} dx \\
&= \left(\frac{17}{15} \right)^p {}_2F_1 \left(-p, 1, \frac{1}{\alpha} + 1; \frac{15}{17} \right).
\end{aligned}$$

Using the fact that $\mathcal{A}^q + \mathcal{B}^q \leq (\mathcal{A} + \mathcal{B})^q$ for $\mathcal{A} \geq 0, \mathcal{B} \geq 0$ with $q \geq 1$, (3.10) gives

$$\begin{aligned}
& \left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] \left(\mathcal{J}(a, b; \varphi) \right)^{-\frac{4\alpha-1}{(b-a)^\alpha}} \right| \\
& \leq \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\ln \varphi^* (a) + \ln \varphi^* \left(\frac{3a+b}{4} \right) \right) \right)
\end{aligned}$$

$$\begin{aligned}
 & \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\ln \varphi^* \left(\frac{3a+b}{4} \right) + \ln \varphi^* \left(\frac{a+b}{2} \right) \right) \right) \\
 & \times \exp \left(\frac{17(b-a)}{240} \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\ln \varphi^* \left(\frac{a+b}{2} \right) + \ln \varphi^* \left(\frac{a+3b}{4} \right) \right) \right) \\
 & \times \exp \left(\frac{b-a}{16} \left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\ln \varphi^* \left(\frac{a+3b}{4} \right) + \ln \varphi^* (b) \right) \right) \\
 & = \left[\left(\varphi^* (a) \varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \varphi^* (b) \right)^{\left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}}} \right. \\
 & \quad \left. \times \left(\varphi^* \left(\frac{3a+b}{4} \right) \left(\varphi^* \left(\frac{a+b}{2} \right) \right)^2 \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\frac{17}{15} \right) \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}}} \right]^{\frac{b-a}{16}}.
 \end{aligned}$$

This completes the proof. $\square$

Corollary 3.7. In Theorem 3.6, if we take $s = 1$, then we get

$$\begin{aligned}
 & \left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] \left(\mathcal{J}(a, b; \varphi) \right)^{-\frac{4\alpha-1}{(b-a)\alpha}} \right| \\
 & \leq \left[\left(\varphi^* (a) \varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \varphi^* (b) \right)^{\left(\frac{1}{\alpha p+1} \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{1}{q}}} \right. \\
 & \quad \left. \times \left(\varphi^* \left(\frac{3a+b}{4} \right) \left(\varphi^* \left(\frac{a+b}{2} \right) \right)^2 \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{1}{2} \right)^{\frac{1}{q}} \left(\frac{17}{15} \right) \left({}_2F_1 \left(-p, 1, \frac{\alpha+1}{\alpha}; \frac{15}{17} \right) \right)^{\frac{1}{p}}} \right]^{\frac{b-a}{16}},
 \end{aligned}$$

where $\mathcal{J}$ is defined as (3.1).

Corollary 3.8. If we take $\alpha = 1$ in Theorem 3.6, we get

$$\begin{aligned}
 & \left| \left[\left(\varphi \left(\frac{3a+b}{4} \right) \right)^{\frac{8}{15}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{-\frac{1}{15}} \left(\varphi \left(\frac{a+3b}{4} \right) \right)^{\frac{8}{15}} \right] \left(\int_a^b (\varphi(x))^{dx} \right)^{\frac{1}{a-b}} \right| \\
 & \leq \left[\left(\varphi^* (a) \varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \varphi^* (b) \right)^{\left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{s+1} \right)^{\frac{1}{q}}} \right. \\
 & \quad \left. \times \left(\varphi^* \left(\frac{3a+b}{4} \right) \left(\varphi^* \left(\frac{a+b}{2} \right) \right)^2 \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{1}{s+1} \right)^{\frac{1}{q}} \left(\frac{17p+1-2p+1}{15p+1(p+1)} \right)^{\frac{1}{p}}} \right]^{\frac{b-a}{16}}.
 \end{aligned}$$

Corollary 3.9. By setting $\alpha = s = 1$ in Theorem 3.6, we get

$$\left| \left(\varphi(x) \varphi(a+b-x) \right)^{\frac{2(x-a)+\rho(a+b-2x)}{2(b-a)}} \left(\varphi \left(\frac{a+b}{2} \right) \right)^{\frac{(1-\rho)(a+b-2x)}{b-a}} \left(\int_a^b (\varphi(x))^{dx} \right)^{\frac{1}{a-b}} \right|$$

$$\leq \left[\left(\varphi^* (a) \varphi^* \left(\frac{3a+b}{4} \right) \varphi^* \left(\frac{a+3b}{4} \right) \varphi^* (b) \right)^{\left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{1}{q}}} \right. \\ \left. \times \left(\varphi^* \left(\frac{3a+b}{4} \right) \left(\varphi^* \left(\frac{a+b}{2} \right) \right)^2 \varphi^* \left(\frac{a+3b}{4} \right) \right)^{\left(\frac{1}{2} \right)^{\frac{1}{q}} \left(\frac{17p+1-2p+1}{15p+1(p+1)} \right)^{\frac{1}{p}}} \right]^{\frac{b-a}{16}}.$$

4. Applications to special means

Let us consider the following means for arbitrary real numbers a, b

The Arithmetic mean: $A(a_1, a_2, \dots, a_n) = \frac{a_1 + a_2 + \dots + a_n}{n}$.

The Harmonic mean: $H(a_1, a_2, \dots, a_n) = \frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}}$

The logarithmic means: $L(a, b) = \frac{b-a}{\ln b - \ln a}$, $a, b > 0$ and $a \neq b$.

The p -Logarithmic mean: $L_p(a, b) = \left(\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right)^{\frac{1}{p}}$, $a, b > 0, a \neq b$ and $p \in \mathbb{R} \setminus \{-1, 0\}$.

Proposition 4.1. Let $a, b \in \mathbb{R}$ with $0 < a < b$, then we have

$$e^{\frac{A^8(a, a, a, b) + A^{-1}(a, b) + A^8(a, b, b, b) - 15L_2^2(a, b)}{15}} \leq e^{\frac{34(a+b)}{15}}.$$

Proof. The assertion follows from Corollary 3.5 applied to the function $\varphi(t) = e^{t^2}$ whose

$$\varphi^*(t) = e^{2t} \text{ and } \left(\int_a^b \varphi(u) du \right)^{\frac{1}{a-b}} = \exp \{ -L_2^2(a, b) \}. \quad \square$$

Proposition 4.2. Let $a, b \in \mathbb{R}$ with $0 < a < b$ and $n > 0$, then we have

$$H^{-\frac{8}{15}}(a, b, b, b) H^{-\frac{1}{15}}(a, b) H^{-15}(a, a, a, b) a^{\frac{nb}{a-b}} b^{-\frac{na}{a-b}} e^{-n} \leq e^{-\left(\frac{5(a+b)^2 + 28ab}{2(a+b)} + \frac{176ab(a+b)}{(3a+b)(a+3b)} \right) \frac{n}{15}}.$$

Proof. The assertion follows from Corollary 3.5 on the interval $[\frac{1}{b}, \frac{1}{a}]$ applied to the func-

$$\text{tion } \varphi(t) = \frac{1}{t^n} \text{ whose } \varphi^*(t) = e^{-\frac{n}{t}} \text{ and } \left(\int_{\frac{1}{b}}^{\frac{1}{a}} \varphi(u) du \right)^{\frac{ab}{a-b}} = a^{\frac{nb}{a-b}} b^{-\frac{na}{a-b}} e^{-n}. \quad \square$$

5. Conclusion

This paper explores inequalities of the corrected dual Simpson-type for fractional multiplicative integrals using multiplicative s -convexity derived from a novel generalized identity. The findings hold the potential to inspire further investigations in this intriguing domain and extensions to other forms of convexity.

Conflict of interest The authors declare that they do not have any conflicts of interest.

References

- [1] Abdeljawad, T., & Grossman, M. (2016). *On geometric fractional calculus*. Journal of Semigroup Theory and Applications, 2016 (2016), Article ID 2, 14 pages.
- [2] Ali, M. A., Abbas, M., Zhang, Z., Sial, I. B., & Arif, R. (2019). *On integral inequalities for product and quotient of two multiplicatively convex functions*. Asian Research Journal of Mathematics, **12**(3), 1-11.
- [3] Ali, M. A., Abbas, M., & Zafer, A. A. (2019). *On some Hermite-Hadamard integral inequalities in multiplicative calculus*. Journal of Inequalities and Special Functions, **10**(1), 111-122. <http://hdl.handle.net/2263/77927>
- [4] Ali, M. A., Budak, H., Sarikaya, M. Z., & Zhang, Z. (2021). *Ostrowski and Simpson type inequalities for multiplicative integrals*. Proyecciones, **40**(3), 743-763. <http://dx.doi.org/10.22199/issn.0717-6279-4136>
- [5] Bashirov, A. E., Kurpinar, E. M., & Özyapici, A. (2008). *Multiplicative calculus and its applications*. Journal of Mathematical Analysis and Applications, **337**(1), 36-48.
- [6] Bashirov, A. E., Misirli, E., & Tandoğdu, Y. (2011). *On modeling with multiplicative differential equations*. Applied Mathematics Journal of Chinese Universities, **26**, 425-438. <https://doi.org/10.1007/s11766-011-2767-6>
- [7] Berhail, A., & Meftah, B. (2022). *Midpoint and trapezoid type inequalities for multiplicatively convex functions*. arXiv preprint arXiv:2208.00535.
- [8] Boulares, H., Meftah, B., Moumen, A., Shafqat, R., Saber, H., Alraqad, T., & Ahmad, E. (2023). *Fractional multiplicative Bullen type inequalities for multiplicative differentiable functions*. Symmetry, **15**(2), 451.
- [9] Budak, H., & Özçelik, K. (2020). *On Hermite-Hadamard type inequalities for multiplicative fractional integrals*. Miskolc Mathematical Notes, **21**(1), 91-99.
- [10] Chiheb, T., Boumaza, N., & Meftah, B. (2020). *Some new Simpson-like type inequalities via prequasiinvexity*. Transylvanian Journal of Mathematics and Mechanics, **12**(1), 1-10.
- [11] Florack, L., Assen, H. V. (2012). *Multiplicative calculus in biomedical image analysis*. Journal of Mathematical Imaging and Vision, **42**, 64-75. <https://doi.org/10.1007/s10851-011-0275-1>
- [12] Fu, H., Peng, Y., & Du, T. (2021). *Some inequalities for multiplicative tempered fractional integrals involving the σ -incomplete gamma functions*. AIMS Mathematics, **6**(7), 7456-7478.
- [13] Grossman, M., & Katz, R. (1972). "Non-Newtonian calculus". Lee Press, Pigeon Cove, Mass.
- [14] Kashuri, A., Meftah, B., Mohammed, P. O., Lupa, A. A., Abdalla, B., Hamed, Y. S., & Abdeljawad, T. (2021). *Fractional weighted Ostrowski type inequalities and their applications*. Symmetry, **13**, 968. <https://doi.org/10.3390/sym13060968>
- [15] Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2006). "Theory and applications of fractional differential equations". North-Holland Mathematics Studies, 204, Elsevier Science B.V., Amsterdam.
- [16] Lakhdari, A., Saleh, W., Meftah, B., & Iqbal, A. (2022). *Corrected dual Simpson type inequalities for differentiable generalized convex functions on a fractal set*. Fractal Fract., **6**(12), 710. <https://doi.org/10.3390/fractalfract6120710>
- [17] Meftah, B., & Azaizia, A. (2019). *Fractional Ostrowski type inequalities for functions whose first derivatives are MT-preinvex*. Revista De Matemáticas De la Universidad del Atlántico Páginas, **6**(1), 33-43.
- [18] Meftah, B. (2023). *Maclaurin type inequalities for multiplicatively convex functions*. Proceedings of the American Mathematical Society, **151**(5), 2115-2125.
- [19] Meftah, B., Lakhdari, A., & Benchettah, D. C. (2022). *Some new Hermite-Hadamard type integral inequalities for twice differentiable s-convex functions*. Computational Mathematics and Modeling, **33**(3), 330-353. <https://doi.org/10.1007/s10598-023-09576-3>
- [20] Meftah, B., & Lakhdari, A. (2023). *Dual Simpson type inequalities for multiplicatively convex functions*. Filomat, **37**(22), 7673-7683.
- [21] Moumen, A., Boulares, H., Meftah, B., Shafqat, R., Alraqad, T., Ali, E. E., & Khaled, Z. (2023). *Multiplicatively Simpson Type Inequalities via Fractional Integral*. Symmetry, **15**(2), 460.
- [22] Pečarić, J. E., Proschan, F., & Tong, Y. L. (1992). "Convex functions, partial orderings, and statistical applications". Mathematics in Science and Engineering, 187. Academic Press, Inc., Boston, MA.
- [23] Peng, Y., Fu, H., & Du, T. (2022). *Estimations of bounds on the multiplicative fractional integral inequalities having exponential kernels*. Communications in Mathematics and Statistics, 1-25.
- [24] Peng, Y., & Du, T. (2023). *Fractional Maclaurin-type inequalities for multiplicatively convex functions and multiplicatively P-functions*. Filomat, **37**(28), 9497-9509.

- [25] Saleh, W., Meftah, B., & Lakhdari, A. (2023). *Quantum dual Simpson type inequalities for q -differentiable convex functions*. International Journal of Nonlinear Analysis and Applications. DOI:10.22075/IJNAA.2023.29280.4109
- [26] Saleh, W., Lakhdari, A., Almutairi, O., & Kiliçman, A. (2023). *Some Remarks on Local Fractional Integral Inequalities Involving Mittag-Leffler Kernel Using Generalized (E, h) -Convexity*. Mathematics, 11(6), 1373. <https://doi.org/10.3390/math11061373>
- [27] Saleh, W., Lakhdari, A., Kiliçman, A., Frioui, A., & Meftah, B. (2023). *Some new fractional Hermite-Hadamard type inequalities for functions with co-ordinated extended (s, m) -prequasiinvex mixed partial derivatives*. Alexandria Engineering Journal, 72, 261-267. <https://doi.org/10.1016/j.aej.2023.03.080>
- [28] Saleh, W., Lakhdari, A., Abdeljawad, T., & Meftah, B. (2023). *On fractional biparameterized Newton-type inequalities*. Journal of Inequalities and Applications, 2023 (1), 1-18.
- [29] Singh, G., & Bhalla, S. (2023). *Two-step Newton's method with multiplicative calculus to solve nonlinear equations*. Journal of Computational Analysis and Applications, 31, 171-179.
- [30] Tan, P., & Du, T. (2023). *On the multi-parameterized inequalities involving the tempered fractional integral operators*. Filomat, 37(15), 4919-4941.
- [31] Ullah, M. I., Ain, Q. T., Khan, A., Abdeljawad, T., & Alqudah, M. A. (2023). *A fractional approach to the solar heating model using an extended ODE system*. Alexandria Engineering Journal, 81, 405-418. <https://doi.org/10.1016/j.aej.2023.09.036>
- [32] Xi, B.-Y., & Qi, F. (2015). *Some integral inequalities of Hermite-Hadamard type for s -logarithmically convex functions*. Acta Mathematica Scientia, English Series, 35A(3), 515-526.
- [33] Xie, J., Ali, M. A., & Sitthiwirattam, T. (2023). *Some new midpoint and trapezoidal type inequalities in multiplicative calculus with applications*. Filomat, 37 (20), 6665-6675.
- [34] Yadav, P., Jahan, S., Shah, K., Peter, O. J., & Abdeljawad, T. (2023). *Fractional-order modeling and analysis of diabetes mellitus: Utilizing the Atangana-Baleanu Caputo (ABC) operator*. Alexandria Engineering Journal, 81, 200-209.
- [35] Yousef, A., Bozkurt, F., Abdeljawad, T., & Emreizeeq, E. (2023). *A mathematical model of COVID-19 and the multi-fears of the community during the epidemiological stage*. Journal of Computational and Applied Mathematics, 419, 114624. <https://doi.org/10.1016/j.cam.2022.114624>