A Common Coincidence of Fixed Point for Generalized Caristi Fixed Point Theorem

JAYASHREE PATIL^a, BASEL HARDAN^{b,*} , AMOL BACHHAV^c

^a Department of Mathematics, Vasantrya Naik Mahavidyalaya, Cidco, Aurangabad, 431003 India.

^{b,*} Department of Mathematics, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431004, India.

^c Navin Jindal School of Management, University of Texas at Dallas, Dallas, 75080, USA.

• Received: 22 February 2021 • Accepted: 24 March 2021 • Published Online: 28 March 2021

Abstract

In this paper, the interpolative Caristi type weakly compatible contractive in a complete metric space is applied to show some common fixed points results related to such mappings. Our application shows that the function which is used to prove the obtained results is a bounded map. An example is provided to show the useability of the acquired results.

Keywords: semi lower continuous, fixed point theorem, Caristi fixed point theorem, contractive mapping.
2010 MSC: 55M20, 37C25, 47H09, 47H10.

1. Introduction and Preliminaries

In 1997, there shown at least five papers that were concerned with fixed point theorems (FPTs) for lower semi-continuous (LSC) multi maps with convex values; see [1, 2, 3, 4, 5]. Most of those theorems are based on one of the well-known selection theorems, for example, [6] and some have applications to the existence of equilibrium points of abstract economies. Our aim in this paper is to derive coincide and common FPTs for LSC multi maps by contractive weakly compatible mappings in Banach metric space. Several studies have extended the Banach contraction principle (PCB) since its inception, from those recent studies we refer to [7, 8, 9, 10]. This paper is inspired by some new work on the extension of the Banach Contraction Principle (BCP) to metric spaces with a partial order [11]. Caristi's fixed point theorem (CFPT) is possibly one of the nicest extensions of the

*Corresponding author: bassil2003@gmail.com

BCP [12, 13]. This theorem states that a map $F : X \rightarrow X$ has a fixed point, where X is a complete metric space and there exists an LSC map $\psi : X \rightarrow [0, +\infty)$ such that

$$d(x, Fx) \leq \psi(x) - \psi(Fx), \quad \forall x \in X. \quad (1.1)$$

A fixed point of F if $F(x) = x$, $\forall x \in X$. In general, the fixed point theorems have established many applications in nonlinear analysis. For example, this result yields from basically all the known natural results of geometrical fixed point theorems in Banach spaces which using the proofing approaching Caristi's theorem by various styles (see [14, 15, 16]). However, it is worth remembering that because of Caristi's theorem's close link to Ekeland's [17] variational concept, many researchers indicate it as Caristi-Ekeland's FPT. For information on Ekeland's variational concept, and equality between Caristi-Ekeland's FPT and the complete metric spaces, see [18]. The purpose of this paper is to present a new generalization of CFPT on the BCP.

Definition 1.1. [19] The LSC map at x_* in a metric space X can be expressed as

$$\lim_{x \rightarrow x_*} \sup \psi(x) \leq \psi(x_*), \quad \forall x_*, x \in X. \quad (1.2)$$

The following result was introduced by Caristie in [12] as:

Theorem 1.2. *Let X be a complete metric space and $\psi : X \rightarrow \mathbb{R}^+$ be an LSC and bonded below function, Suppose that F is a Caristie type mapping on X dominated by ψ . If F satisfies*

$$d(x, Fx) \leq \psi(x) - \psi(Fx), \quad \forall x \in X. \quad (1.3)$$

Then, F has a fixed point in X .

Definition 1.3. [20] Let F and T are selfing maps in X , then

- (i) A point $x_0 \in X$ is said to be a common fixed point of F and T if $x_0 = Fx_0 = Tx_0$.
- (ii) A point $x_0 \in X$ is called a coincidence point of F and T if $x_0 = Fx = Tx$.
- (iii) The mappings $F, T : X \rightarrow X$ are said to be weakly compatible if $FTx = TFx$ whenever $Fx = Tx$.

Lemma 1.4. [20] *Let X be a non-empty set and suppose that the mappings $F, T : X \rightarrow X$ have a unique coincidence point $x_0 \in X$. If F and T are weakly compatible, then F and T have a unique common fixed point.*

Du in [19] has recalled another LSC function to get a direct proof of CFPT. Like this work depends on some development in BCP in metric space, see [21, 22, 23, 24]. Such these studies are still being replenished. A lot of recent papers study fixed point theorem by Caristi's ways. Among these recent studies we will point out to [25, 26, 27, 28, 29, 30, 31]. Here, we use a LSC function to obtain some new FPTs of Caristi type for a weakly compatible contractive mapping in the Banach metric space.

2. Main results

In this section, we will prove the uniqueness of coincidence and a common fixed point in Banach space for generalization CFPT type.

Theorem 2.1. *Let F, T are weakly compatible contractive self-mappings on a Banach space X , and let $\psi : X \rightarrow \mathbb{R}^+$ be a LSC function such that,*

$$\psi x \leq d(Fx, Tx), \quad (2.1)$$

for all $x \in X$, satisfies

$$d(Fx, Fy) \leq \psi(Tx) - \psi(Fx) + \psi(Ty) - \psi(Fy), \quad (2.2)$$

for all $x, y \in X$ and $x \neq y$. Then F and T have a unique common fixed point.

Proof. First: Existence.

Consider $Fx_{n-1} = Tx_n = x_n$ for all $n \in \mathbb{N}$. Now,

$$\begin{aligned} d(x_n, x_{n+1}) &= d(Fx_{n-1}, Fx_n) \leq \psi(Tx_{n-1}) - \psi(Fx_{n-1}) + \psi(Tx_n) - \psi(Fx_n) \\ &= \psi(x_{n-1}) - \psi(x_n) + \psi(x_n) - \psi(x_{n+1}) \\ &= \psi(x_{n-1}) - \psi(x_{n+1}) \\ &\leq d(Fx_{n-1}, Tx_{n-1}) - d(Fx_{n+1}, Tx_{n+1}). \end{aligned}$$

We obtain,

$$d(x_n, x_{n+2}) \leq d(x_{n-1}, x_n) \quad (2.3)$$

Since F, T are contractive mappings then they have continuous property, by Gopal in [32], the continuity and down bounded limit for ψ lead us to, x_n is decreasing to some points $x_0 \in X$, for all $x \in X$

$$\psi(x_0) \leq \psi(x).$$

Since X is a Banach space thus, $X \supset \{x_n\} \rightarrow x_0$. Also, $\{x_n\} \supset \{x_{n_k}\} \rightarrow x_0$ for all $n, k \in \mathbb{N}$, we have

$$\lim_{n, k \rightarrow \infty} x_{n_k} = F(\lim_{n, k \rightarrow \infty} x_{n_k}) = F(x_0), \quad (2.4)$$

$$\lim_{n \rightarrow \infty} x_n = T(\lim_{n \rightarrow \infty} x_n) = T(x_0). \quad (2.5)$$

Hence, x_0 is a coincidence fixed point of F and T and by Lemma 1.4, we obtain that x_0 is a common fixed point of F and T . Existence has been proven.

Second: Uniqueness.

Suppose that $F(x_*) = T(x_*) = x_*$ for all $x_* \in X$ such that $x_0 \neq x_*$, so by (2.2) we get

$$\begin{aligned} d(x_*, x_0) &= d(Fx_*, Fx_0) \\ &\leq \psi(Tx_*) - \psi(Fx_*) + \psi(Tx_0) - \psi(Fx_0) \\ &\leq \psi x_* - \psi x_* + \psi x_0 - \psi x_0 \\ &\leq d(Fx_*, Tx_*) - d(Fx_*, Tx_*) + d(Fx_0, Tx_0) - d(Fx_0, Tx_0) \\ &\leq 0. \end{aligned}$$

Hence, $x_* = x_0$. Uniqueness has been proven. $\square$

We will generalize Theorem 2.1 for all repetitions $m \in \mathbb{N}$.

Theorem 2.2. Let F^m, T^m , $m \in \mathbb{N}$ are weakly compatible contractive self-maps on Banach space X , and let $\psi : X \rightarrow \mathbb{R}^+$ be a SLC function such that,

$$\psi x \leq d(F^m x, T^m x), \quad (2.6)$$

for all $x \in X$, satisfies

$$d(F^m x, F^m y) \leq \psi(T^m x) - \psi(F^m x) + \psi(T^m y) - \psi(F^m y), \quad (2.7)$$

for all $x, y \in X$ and $x \neq y$. Then F and T have a unique common fixed point.

Proof. First: Existence.

F^m and T^m are continuous maps, since F^m and T^m are contractive. In the same way as proof Theorem 2.1, we conclude that F^m and T^m have a unique fixed common point in X . Suppose $F^m x_0 = T^m x_0 = x_0$. Let $F^r x_0 \neq T^r x_0 \neq x_0$ for $r = 1, 2, \dots, m-1$. Also, let $F^r x_0 = T^r x_0 = x_*$ for all $x_0 \neq x_*$ then

$$\begin{aligned} d(F^r x_0, F^m x_0) &= d(F F^{r-1} x_0, F F^{m-1} x_0) \\ &\leq \psi(T F^{r-1} x_0) - \psi(F F^{r-1} x_0) + \psi(T F^{m-1} x_0) - \psi(F F^{m-1} x_0) \\ &\leq \psi(T F^{r-1} x_0) - \psi(F^r x_0) + \psi(T F^{m-1} x_0) - \psi(F^m x_0) \\ &= 0, \quad m > 1. \end{aligned}$$

Therefore, $F^r x_0 = T^r x_0 = F^m x_0 = x_0$, for all $r = 1, 2, \dots, m-1$. Thus, $F x_0 = T x_0 = x_0$, and since F and T are weakly compatible maps. Hence, x_0 is a common fixed point for F and T .

Second: Uniqueness

Proving the uniqueness is the same steps as proving the uniqueness of the theorem 2.1. $\square$

In the last section, we shall introduce an application and example to justify the viability and usability of our results.

3. application

Theorem 3.1. A function ψ which is defined in (2.6) can be considered as a Lipschitz bounded function $\forall x \in X$ and $\forall m \in \mathbb{N}$.

Proof. Since, $\psi x \leq d(F^n x, T^n x)$, $\forall n \in \mathbb{N}$. We have for all $x_1, x_2 \in X$ such that $x_1 \neq x_2$.

$$\begin{aligned} \psi x_1 - \psi x_2 &\leq d(F^n x_1, T^n x_1) - d(F^n x_2, T^n x_2) \\ &= d(F^n x_1, T^n x_1) - d(F^n x_2, T^n x_2) + d(F^n x_1, T^n x_2) - d(F^n x_1, T^n x_2) \\ &= d(F^n x_1, T^n x_1) - d(F^n x_1, T^n x_2) + d(F^n x_1, T^n x_2) - d(F^n x_2, T^n x_2) \\ &= d(T^n x_1, T^n x_2) + d(F^n x_1, F^n x_2) \\ &= d(x_1, x_2) + d(F^n x_1, F^n x_2) \\ &\leq M d(x_1, x_2), \quad M \in \mathbb{N}. \end{aligned} \quad (3.1)$$

Then, $M \geq 0$ where, $\psi : X \rightarrow \mathbb{R}^+$. Now we must prove that $M > 1$, so we will study all possible possibilities of M as following

- i When $M = 0$, then $x_1 = x_2$ which is contradiction.
- ii When $M = 1$, then $d(F^n x_1, F^n x_2) = 0$ which is contradiction since F has a unique fixed point.
- iii When $0 < M < 1$ then, there exist $\mathcal{A}$ from (3.1) satisfies

$$d(F^n x_1, F^n x_2) = \mathcal{A}d(x_1, x_2). \quad (3.2)$$

Inequalities (3.1) and (3.2) show that $\mathcal{A}$ is negative real number, which is a contradiction.

Therefore, $\mathcal{A} > 1$. Hence, ψ is bounded Lipschitz function. $\square$

Example 3.2. Consider, $X = [0, 1)$. Take $\psi : [0, 1) \rightarrow \mathbb{R}^+$ defined by

$$\psi u = \begin{cases} 0, & \text{if } u \geq 0, \\ 1, & \text{if } u < 1. \end{cases}$$

Suppose, $F(u) = u^2$ and $T(u) = u$. Then, for all $u, v \in X$ we get

$$\begin{aligned} d(F(u), F(v)) &= d(u^2, v^2) \\ &< d(u, v). \end{aligned} \quad (3.3)$$

Clear that F is a contractive mapping. By inequalities (2.1) and (2.2) we get

$$\begin{aligned} d(Fu, Fv) &\leq d(F(Tu), T^2u) - d(F^2u, T(Fu)) + d(F(Tv), T^2v) - d(F^2v, T(Fv)) \\ &= d(u^2, u) - d(u^4, u^2) + d(v^2, v) - d(v^4, v^2) \end{aligned} \quad (3.4)$$

$$\leq d(u^4, u) + d(v^4, v). \quad (3.5)$$

We have two possibilities,

First possibility: If $u = v$, then F and T have a coincidence fixed point, and since F and T are commutable mappings then F and T have a unique common fixed point which is $u = 0 \in X$.

Second possibility: If $u \neq v$, then (3.5) is satisfies for every $u, v \in X$.

4. conclusion

In this work, we have obtained the uniqueness of some common fixed points by utilized the interpolative Caristi type weakly compatible contractive results concerning such mappings through the Banach metric space. On other hand, Bakhtin in [33] introduced the concept of a quasi-metric space and showed some fixed point theorems on such space. Recently, Kamran in [34] generalized the quasi metric space and showed new primary notions. The problem considered in this paper can be studied on quasi metric space, which is a direction we are working on.

Acknowledgement

We would like to thank the referees very much for their valuable comments and suggestions.

References

- [1] Kim W, Lee K (1997), *Existence of equilibrium and separation in generalized games*, J. Math. Anal. Appl. **207**(2): 316–325.
- [2] O'Regan D (1997), *Fixed point theory for compact upper semi-continuous or lower semi-continuous set valued maps*, Proc. Amer. Math. Soc. **125**(3): 875–881.
- [3] Wu X (1997), *A new fixed point theorem and its applications*, Proc. Amer. Math. Soc. **125**: 1779–1783.
- [4] Zhang X (1997), *Some intersection theorems and minimax inequalities*, J. Optim. Th. and Appl. **94**: 195–207.
- [5] Zheng X (1997), *Approximate selection theorems and their applications*, J. Math. Anal. Appl. **212**: 88–97.
- [6] Michael E (1966), *A selection theorem*, Proc. Amer. Math. Soc. **17**: 1404–1406.
- [7] Patil J, Hardan B (2019), *On fixed point theorems in complete metric space*, Journal of computer and mathematical Sciences, **10**(7): 1419–1425. [https:// DOI: 10.29055/jcms/1128](https://doi.org/10.29055/jcms/1128)
- [8] Patil J, Hardan B, Abdo M S, Chaudhari A, Bachhav A (2020), *A fixed point theorem for Hardy-Rogers type on generalized fractional differential equations*, Advances in the Theory of Nonlinear Analysis and its Applications, **4**(4): 407–420. [https://DOI: 10.31197/atnaa.767331](https://doi.org/10.31197/atnaa.767331)
- [9] Patil J, Hardan B, Chaudhari A, Bachhav A (2020), *Approximate fixed points for n -Linear functional by (μ, σ) - nonexpansive Mappings on n -Banach spaces*, Journal of Mathematical Analysis and Modeling, **1**(1): 20-32. [https:// DOI: 10.48185/jmam.v1i1.23](https://doi.org/10.48185/jmam.v1i1.23)
- [10] 2) Patil J, Hardan B, Abdo M S, Chaudhari A, Bachhav A (2021), *Generalized Fractional Differential Equations by Using a Fixed Point Theorem for Generalized Contractive Type*, Dynamics of Continuous, Discrete and Impulsive Systems, **28**(2), (2021), 77-88.
- [11] Nieto J, Rodriguez-Lopez R, *Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations*, Order. **22**(3): 223–239. [https:// DOI:10.1007/s11083-005-9018-5](https://doi.org/10.1007/s11083-005-9018-5)
- [12] Caristi J (1976), *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc. **215**: 241-251.
- [13] Caristi J (1979), *Fixed point theory and inwardness conditions*, Applied Nonlinear Analysis , 479–483.
- [14] Brøndsted A (1976), *Fixed point and partial orders*, Proc. Amer. Math. Soc. **60**: 365–366.
- [15] Browder F (1975), *On a theorem of Caristi and Kirk*, Proc. Seminar on Fixed Point Theory and Its Applications, Dalhousie University, Academic Press, 23–27.
- [16] Kirk W, Caristi J (1975), *Mapping Theorems in Metric and Banach Spaces*, Bull. L'Acad. Polon. Sci. **25**:891–894.
- [17] Ekeland I (1972), *Sur les problemes variationnels*, Comptes Rendus Acad. Sci. Paris, **275**: 1057–1059.
- [18] Sullivan F (1981), *A Characterization of Complete Metric Spaces*, Proc. Amer. Math. Soc., **83**(2): 345–346.
- [19] Due W (2016), *A direct proof of Caristi's fixed point Theorem*, Applied Mathematical Sciences, **10**(46): 2289–2294. <http://dx.doi.org/10.12988/ams.2016.66190>
- [20] Arshad M, Ahmad J, Karapinar E (2013), *Some Common Fixed Point Results in Rectangular Metric Spaces*, International Journal of Analysis **2013**(2013):1–7. <https://doi.org/10.1155/2013/307234>
- [21] Branciari A (2000), *A fixed point theorem of Banach-Caccioppoli type on a class of generalized metric spaces*, Publicationes Mathematicae Debrecen **57**(1-2): 31–37.
- [22] Cho S (2018), *Fixed Point Theorems for generalized weakly contraction mappings in metric space with applications*, Fixed point Theorem and Applications springer open Journal, **3**: 1–18. <https://doi.org/10.1186/s13663-018-0628-1>
- [23] Patil S, Dolhar U (2016), *A note on Development of metric fixed point theory*, Int. J. Adv. Res. **8**(4): 1729–1734. <http://dx.doi.org/10.21474/IJAR01/1384>
- [24] Suzuki T (2014), *Generalized metric spaces do not have the compatible Topology*, Hindawi Publishing Corporation, Abstract and Applied Analysis, **2014**: 5 pages. <https://doi.org/10.1155/2014/458098>
- [25] Aamri M, Chaira K, Lazaiz S, Marhrani El-M (2020), *Caristi type fixed point theorems using Szász principle in quasi-metric spaces*, Carpathian Journal of Mathematics, **36**(2): 179–188.

- [26] Altun I, Sahin H, Turkoglu D (2020), *Caristi-type fixed point theorems and some generalizations on M-metric space*, Bulletin of the Malaysian Mathematical Sciences Society, **43**(3): 2647–2657. <https://doi.org/10.1007/s40840-019-00823-8>
- [27] Aslantas M, Sahin H, Turkoglu D (2020), Some Caristi type fixed point theorems, The Journal of Analysis, 1–15. <https://doi.org/10.1007/s41478-020-00248-8>
- [28] Chuensupantharat N, Gopal D (2020), *On Caristi's fixed point theorem in metric spaces with a graph*, Carpathian Journal of Mathematics, **36**(2): 259–268.
- [29] Kuhlmann F, Kuhlmann K (2018), Paulsen M, *The Caristi–Kirk Fixed Point Theorem from the point of view of ball spaces*, Journal of Fixed Point Theory and Applications, **20**: p. 107. <https://doi.org/10.1007/s11784-018-0576-8>
- [30] Pant A, Pant R, Joshi M (2019), Caristi type and meir-keeler type fixed point theorems, Filomat, **33**(12): 3711–3721. <https://doi.org/10.2298/FIL1912711P>
- [31] Patil J, Hardan B, Chaudhari A, Bachhav A (2020), *Caristi type fixed point theorems of contractive mapping with application*, One day national conference on recent advances in sciences, Commerce and science college Kannad, Aurangabad, **38**: 609–614.
- [32] Gopal D, Kumar D, Sukla S, Banach Fixed Point Theorem and its Generalizations, Taylor and Francis Group, (2018).
- [33] Bakhtin I. A (1989), *The contraction mapping principle in almost metric spaces*, Funct. Anal., **30**: 26–37.
- [34] Kamran T, Samreen M, UL Ain Q (2017), *A Generalization of b-Metric Space and Some Fixed Point Theorems*, Mathematics, **5**(2): 19 pages. <http://doi:10.3390/math5020019>